Mathematics Applications Units 1 and 2 Formula Sheet

Measurement

Pythagoras' theorem

$$a^2 + b^2 = c^2$$

Circumference of circle, radius r

$$C = 2\pi r$$

Arc length of circle, central angle θ

$$l = \frac{\theta}{360} 2\pi r$$

Area of circle

$$A = \pi r^2$$

Area of sector

$$A = \frac{\theta}{360}\pi r^2$$

Area of triangle, base b and height h

$$A = \frac{1}{2}bh$$

Area of parallelogram

$$A = bh$$

Area of trapezium, parallel sides a and b

$$A = \frac{1}{2}(a+b)h$$

Volume of prism, base area A

$$V = Ah$$

Volume of pyramid

$$V = \frac{1}{3}Ah$$

Surface area of cylinder

$$S = 2\pi r h + 2\pi r^2$$

Volume of cylinder

$$V = \pi r^2 h$$

Surface area of cone, slant height s

$$S = \pi r^2 + \pi r s$$

Volume of cone

$$V = \frac{1}{3}\pi r^2 h$$

Surface area of sphere

$$S = 4\pi r^2$$

Volume of sphere

$$V = \frac{4}{3}\pi r^3$$

Trigonometry

In a right triangle:

$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$\tan \theta = \frac{\text{opposite}}{\text{adjacent}}$$

Sine rule

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Cosine rule

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

Area

$$A = \frac{1}{2}ab\sin C$$

$$A = \sqrt{s(s-a)(s-b)(s-c)}$$
 where $s = \frac{a+b+c}{2}$

Linear equations

Gradient

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

Equation

$$y = mx + c$$

Mathematics Applications Units 1 and 2 Formula Sheet

Consumer arithmetic

P is the principal, *A* is the final balance, *I* is the interest, *r* is the annual rate of interest as a decimal, *n* is the number of compounding periods and *t* is the time in years.

Simple interest

$$I = Prt$$

Compound interest

$$A = P\left(1 + \frac{r}{n}\right)^{nt}$$

 $PE = \frac{\text{Market value per share}}{\text{Earnings per share}}$

Univariate data analysis

Standard score, for normally distributed data set with mean \bar{x} and standard deviation s_x

$$z = \frac{x - \bar{x}}{s_x}$$

In a normal distribution, approximately 68% of values lie within one, 95% of values lie within two and 99.7% of values lie within three, standard deviations of the mean.

In a data set with lower quartile Q_1 , upper quartile Q_3 and interquartile range IQR, possible outliers lie below $Q_1-1.5IQR$ or above $Q_3+1.5IQR$.