



Standard Reduction Potentials at 25 °C

Half-reaction	E°(volts)
$F_2(g) + 2 e^- \rightleftharpoons 2 F^-(aq)$	+ 2.87
$H_2O_2(aq) + 2 H^+(aq) + 2 e^- \rightleftharpoons 2 H_2O(l)$	+ 1.78
$PbO_2(s) + SO_4^{2-}(aq) + 4 H^+(aq) + 2 e^- \rightleftharpoons PbSO_4(s) + 2 H_2O(l)$	+ 1.69
$2 HC_2O(aq) + 2 H^+(aq) + 2 e^- \rightleftharpoons C_2O_2(g) + 2 H_2O(l)$	+ 1.63
$MnO_4^-(aq) + 8 H^+(aq) + 5 e^- \rightleftharpoons Mn^{2+}(aq) + 4 H_2O(l)$	+ 1.51
$Au^{3+}(aq) + 3 e^- \rightleftharpoons Au(s)$	+ 1.50
$HC_2O(aq) + H^+(aq) + 2 e^- \rightleftharpoons C_2O^-(aq) + H_2O(l)$	+ 1.49
$PbO_2(s) + 4 H^+(aq) + 2 e^- \rightleftharpoons Pb^{2+}(aq) + 2 H_2O(l)$	+ 1.46
$Cl_2(g) + 2 e^- \rightleftharpoons 2 Cl^-(aq)$	+ 1.36
$Cr_2O_7^{2-}(aq) + 14 H^+(aq) + 6 e^- \rightleftharpoons 2 Cr^{3+}(aq) + 7 H_2O(l)$	+ 1.33
$O_2(g) + 4 H^+(aq) + 4 e^- \rightleftharpoons 2 H_2O(l)$	+ 1.23
$Br_2(l) + 2 e^- \rightleftharpoons 2 Br^-(aq)$	+ 1.07
$Ag^+(aq) + e^- \rightleftharpoons Ag(s)$	+ 0.80
$NO_3^-(aq) + 2 H^+(aq) + e^- \rightleftharpoons NO_2(g) + H_2O(l)$	+ 0.80
$Fe^{3+}(aq) + e^- \rightleftharpoons Fe^{2+}(aq)$	+ 0.77
$O_2(g) + 2 H^+(aq) + 2 e^- \rightleftharpoons H_2O_2(aq)$	+ 0.68
$I_2(s) + 2 e^- \rightleftharpoons 2 I^-(aq)$	+ 0.54
$O_2(g) + 2 H_2O(l) + 4 e^- \rightleftharpoons 4 OH^-(aq)$	+ 0.40
$Cu^{2+}(aq) + 2 e^- \rightleftharpoons Cu(s)$	+ 0.34
$S(s) + 2 H^+(aq) + 2 e^- \rightleftharpoons H_2S(aq)$	+ 0.14
$2 H^+(aq) + 2 e^- \rightleftharpoons H_2(g)$	0 exactly
$Pb^{2+}(aq) + 2 e^- \rightleftharpoons Pb(s)$	- 0.13
$Sn^{2+}(aq) + 2 e^- \rightleftharpoons Sn(s)$	- 0.14
$Ni^{2+}(aq) + 2 e^- \rightleftharpoons Ni(s)$	- 0.25
$Co^{2+}(aq) + 2 e^- \rightleftharpoons Co(s)$	- 0.28
$PbSO_4(s) + 2 e^- \rightleftharpoons Pb(s) + SO_4^{2-}(aq)$	- 0.36
$Cd^{2+}(aq) + 2 e^- \rightleftharpoons Cd(s)$	- 0.40
$2 CO_2(g) + 2 H^+(aq) + 2 e^- \rightleftharpoons HOOCOOH(aq)$	- 0.43
$Fe^{2+}(aq) + 2 e^- \rightleftharpoons Fe(s)$	- 0.44
$Cr^{3+}(aq) + 3 e^- \rightleftharpoons Cr(s)$	- 0.74
$Zn^{2+}(aq) + 2 e^- \rightleftharpoons Zn(s)$	- 0.76
$2 H_2O(l) + 2 e^- \rightleftharpoons H_2(g) + 2 OH^-(aq)$	- 0.83
$Mn^{2+}(aq) + 2 e^- \rightleftharpoons Mn(s)$	- 1.18
$Al^{3+}(aq) + 3 e^- \rightleftharpoons Al(s)$	- 1.66
$Mg^{2+}(aq) + 2 e^- \rightleftharpoons Mg(s)$	- 2.37
$Na^+(aq) + e^- \rightleftharpoons Na(s)$	- 2.71
$Ca^{2+}(aq) + 2 e^- \rightleftharpoons Ca(s)$	- 2.87
$Str^{2+}(aq) + 2 e^- \rightleftharpoons Sr(s)$	- 2.89
$Ba^{2+}(aq) + 2 e^- \rightleftharpoons Ba(s)$	- 2.91
$K^+(aq) + e^- \rightleftharpoons K(s)$	- 2.93

Copyright

© Curriculum Council, 2010

This document—apart from any third party copyright material contained in it—may be freely copied, or communicated on an intranet, for non-commercial purposes by educational institutions, provided that it is not changed in any way and that the Curriculum Council is acknowledged as the copyright owner.

Copying or communication for any other purpose can be done only within the terms of the Copyright Act or by permission of the Curriculum Council.

Copying or communication of any third party copyright material contained in this document can be done only within the terms of the Copyright Act or by permission of the copyright owners.

Periodic Table

1 H 1.008	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18																	
3 Li 6.941	4 Be 9.012	5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18	11 Na 22.99	12 Mg 24.31	13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95	19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.88	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.38	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80	
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc 101.1	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3	55 Cs 132.9	56 Ba 137.3	57 La 138.9	58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm 144.9	62 Sm 150.4	63 Eu 152.0	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0	71 Lu 175.0
87 Fr 226.0	88 Ra 226.0	89 Ac 226.0	90 Th 232.0	91 Pa 232.0	92 U 238.0	93 Np 238.0	94 Pu 238.0	95 Am 238.0	96 Cm 238.0	97 Bk 238.0	98 Cf 238.0	99 Es 238.0	100 Fm 238.0	101 Md 238.0	102 No 238.0	103 Lr 238.0	104 Rf 238.0	105 Db 238.0	106 Sg 238.0	107 Bh 238.0	108 Hs 238.0	109 Mt 238.0	110 Ds 238.0	111 Rg 238.0	112 Og 238.0	113 Nh 238.0	114 Fl 238.0	115 Mc 238.0	116 Lv 238.0	117 Ts 238.0	118 Og 238.0			

6
C
12.01

← Atomic Number
← Symbol
← Atomic Mass

* Lanthanide Series
** Actinide Series

Formulae

Number of moles $n = \frac{m}{M} = \frac{\text{mass}}{\text{molar mass}}$

Number of moles of solute $n = cV$

Number of moles of a gas at STP $n = \frac{V}{22.41}$

Ideal gas law $PV = nRT$

Parts per million $\text{ppm} = \frac{\text{mass of solute (mg)}}{\text{mass of solution (kg)}}$

pH of a solution $\text{pH} = -\log [\text{H}^+]$

Density $\rho = \frac{\text{mass of sample}}{\text{volume of sample}}$

Units

Volumes are given in the units of litres (L), or millilitres (mL).
Temperatures are given in the units of degrees Celsius (°C) or Kelvin (K).
It may be assumed that 0.0°C = 273.1 K
Energy changes are given in the SI unit kilojoule (kJ)
Pressures are given in the SI unit kilopascal (kPa) and in atmospheres (atm).
1.00 atm = 101.3 kPa
Solution concentrations are given in the units moles per litre (mol L⁻¹), grams per litre (g L⁻¹) and parts per million.

Constants

Universal gas constant, R = 8.315 J K⁻¹ mol⁻¹
Avogadro constant, N = 6.022 x 10²³ mol⁻¹
Volume of 1.000 mol of an ideal gas at 0.0°C and 101.3 kPa is 22.41 L
S.T.P. is 0.0°C and 101.3 kPa
Equilibrium constant for water at 25°C, K_w = 1.00 x 10⁻¹⁴

Solubility rules for ionic solids in water

Soluble in water

Soluble	Exceptions	
	Insoluble	Slightly soluble
Most chlorides	AgCl	PbCl ₂
Most bromides	AgBr	PbBr ₂
Most iodides	AgI, PbI ₂	
All nitrates	No exceptions	
All ethanoates	No exceptions	
Most sulfates	SrSO ₄ , BaSO ₄ , PbSO ₄	CaSO ₄ , Ag ₂ SO ₄

Insoluble in water

Insoluble	Exceptions	
	Soluble	Slightly soluble
Most hydroxides	NaOH, KOH, Ba(OH) ₂ (note: NH ₄ OH and AgOH do not exist)	Ca(OH) ₂ , Sr(OH) ₂
Most carbonates	Na ₂ CO ₃ , K ₂ CO ₃ , (NH ₄) ₂ CO ₃	
Most phosphates	Na ₃ PO ₄ , K ₃ PO ₄ , (NH ₄) ₃ PO ₄	
Most sulfides	Na ₂ S, K ₂ S, (NH ₄) ₂ S	

Soluble = more than 0.1 mole dissolves per litre
Slightly soluble = between 0.01 and 0.1 mole dissolves per litre
Insoluble = less than 0.01 mole dissolves per litre

Colours of selected ionic substances

In general, ionic solids have the same colour as that of any coloured ion they contain. Two colourless ions in general produce a white solid. Selected exceptions to these two basic rules are noted below.

Ionic Solid	Colour
calcium iodide	pale yellow
copper(I) carbonate	green
copper(II) chloride	green
copper(II) oxide	black
copper(II) sulfide	black
iron(III) sulfide	black
lead(II) iodide	yellow
lead(II) sulfide	black
manganese(II) sulfide	black
silver carbonate	yellow
silver iodide	yellow
silver oxide	brown/black
silver sulfide	black

Other coloured substances

Most gases and liquids are colourless, and most metals are silvery or grey. Selected exceptions to these basic rules are noted below.

Substance	State	Colour
copper	solid	salmon pink
gold	solid	yellow
nitrogen dioxide	gas	brown
phosphorus	solid	yellow
sulfur	liquid, solid	yellow

Coloured ions in aqueous solution

Cation	Colour
Cr ³⁺	deep green
Co ²⁺	pink
Cu ²⁺	blue
Fe ²⁺	pale green
Fe ³⁺	brown
Mn ²⁺	very pale pink
Ni ²⁺	green

Anion	Colour
CrO ₄ ²⁻	yellow
Cr ₂ O ₇ ²⁻	orange
MnO ₄ ⁻	purple

Coloured halogens

Halogen	Colour of free element
F ₂ (g)	yellow
Cl ₂ (g)	green
Br ₂ (l)	red
I ₂ (s)	dark grey

Halogen	Colour of halogen in aqueous solution
Cl ₂ (aq)	pale yellow
Br ₂ (aq)	orange
I ₂ (aq)	brown

Halogen	Colour of halogen in organic solvent
Br ₂	red
I ₂	purple