
Standard Reduction Potentials at 25 °C

Half-reaction			E°(volts)
F ₂ (g) + 2 e	\rightleftharpoons	2 F ⁻ (aq)	+ 2.89
$H_2O_2(aq) + 2 H^+(aq) + 2 e^{-\frac{1}{4}}$	\rightleftharpoons	2 H ₂ O(<i>l</i>)	+ 1.76
$PbO_2(s) + SO_4^{2-}(aq) + 4 H^+(aq) + 2 e^{-\frac{1}{4}}$	ightharpoonup	$PbSO_4(s) + 2 H_2O(\ell)$	+ 1.69
2 HClO(aq) + 2 H ⁺ (aq) + 2 e	-	$C\ell_2(g) + 2 H_2O(\ell)$	+ 1.63
$MnO_4^-(aq) + 8 H^+(aq) + 5 e^{-\frac{1}{4}}$	\rightleftharpoons	$Mn^{2+}(aq) + 4 H_2O(\ell)$	+ 1.51
Au ³⁺ (aq) + 3 e	\rightleftharpoons	Au(s)	+ 1.50
HC{O(aq) + H ⁺ (aq) + 2 e	-	$C\ell^-(aq) + H_2O(\ell)$	+ 1.49
$PbO_{2}(s) + 4 H^{+}(aq) + 2 e^{-\frac{1}{3}}$	ightharpoonup	$Pb^{2+}(aq) + 2 H_2O(\ell)$	+ 1.46
$C\ell_2(g) + 2 e^{-\frac{\pi}{4}}$	ightharpoonup	2 Cl-(aq)	+ 1.36
$\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14 \text{ H}^+(\text{aq}) + 6 \text{ e}^{-\frac{1}{4}}$	=	$2 \text{ Cr}^{3+}(aq) + 7 \text{ H}_2O(\ell)$	+ 1.36
$O_2(g) + 4 H^+(aq) + 4 e^{-\frac{1}{4}}$	ightharpoonup	2 H ₂ O(<i>l</i>)	+ 1.23
$Br_2(\ell) + 2 e^{-\frac{1}{4}}$	-	2 Br ⁻ (aq)	+ 1.08
Ag ⁺ (aq) + e	\rightleftharpoons	Ag(s)	+ 0.80
Fe ³⁺ (aq) + e ⁻	ightharpoonup	Fe ²⁺ (aq)	+ 0.77
$O_{2}(g) + 2 H^{+}(aq) + 2 e^{-\frac{1}{4}}$	\rightleftharpoons	$H_2O_2(aq)$	+ 0.70
$I_{2}(s) + 2 e^{-\frac{1}{3}}$	\rightleftharpoons	2 I ⁻ (aq)	+ 0.54
$O_2(g) + 2 H_2O(l) + 4 e^{-\frac{1}{2}}$	\rightleftharpoons	4 OH-(aq)	+ 0.40
Cu ²⁺ (aq) + 2 e	=	Cu(s)	+ 0.34
S(s)+ 2 H ⁺ (aq) + 2 e	ightharpoonup	H ₂ S(aq)	+ 0.17
2 H ⁺ (aq) + 2 e	-	$H_2(g)$	0 exactly
Pb ²⁺ (aq) + 2 e	=	Pb(s)	- 0.13
Sn ²⁺ (aq) + 2 e	\rightleftharpoons	Sn(s)	- 0.14
Ni ²⁺ (aq) + 2 e	-	Ni(s)	-0.24
Co ²⁺ (aq) + 2 e	\rightleftharpoons	Co(s)	-0.28
$PbSO_{4}(s) + 2 e^{-\frac{1}{3}}$	ightharpoonup	$Pb(s) + SO_4^{2-}(aq)$	-0.36
$Cd^{2+}(aq) + 2e^{-\frac{1}{4}}$	=	Cd(s)	-0.40
$2 CO_{2}(g) + 2 H^{+}(aq) + 2 e^{-\frac{1}{4}}$	ightharpoonup	$H_2C_2O_4(aq)$	-0.43
Fe ²⁺ (aq) + 2 e	\rightleftharpoons	Fe(s)	-0.44
Cr ³⁺ (aq) + 3 e	-	Cr(s)	-0.74
Zn ²⁺ (aq) + 2 e	\rightleftharpoons	Zn(s)	-0.76
2 H ₂ O(ℓ) + 2 e	\rightleftharpoons	$H_2(g) + 2 OH^-(aq)$	- 0.83
Mn ²⁺ (aq) + 2 e	ightharpoonup	Mn(s)	– 1.18
$A\ell^{3+}(aq) + 3 e^{-\frac{1}{4}}$	ightharpoonup	Al(s)	– 1.68
Mg ²⁺ (aq) + 2 e ⁻ =	-	Mg(s)	- 2.36
Na ⁺ (aq) + e	ightharpoonup	Na(s)	- 2.71
Ca ²⁺ (aq) + 2 e	-	Ca(s)	- 2.87
Sr ²⁺ (aq) + 2 e		. ,	- 2.90
Ba ²⁺ (aq) + 2 e	ightharpoonup	Ba(s)	- 2.91
K ⁺ (aq) + e	-	K(s)	- 2.94

[Data source: Aylward, G.H., & Findlay, T. (2008). SI Chemical Data (6th ed.). Queensland: John Wiley & Sons Australia, Ltd.]





CHEMISTRY DATA SHEET 2013

Copyright

© School Curriculum and Standards Authority, 2012

This document – apart from any third party copyright material contained in it – may be freely copied, or communicated on an intranet, for non-commercial purposes in educational institutions, provided that it is not changed and that the School Curriculum and Standards Authority is acknowledged as the copyright owner, and that the Authority's moral rights are not infringed.

Copying or communication for any other purpose can be done only within the terms of the *Copyright Act 1968* or with prior written permission of the School Curriculum and Standards Authority. Copying or communication of any third party copyright material can be done only within the terms of the *Copyright Act 1968* or with permission of the copyright owners.

Any content in this document that has been derived from the Australian Curriculum may be used under the terms of the <u>Creative Commons Attribution-NonCommercial 3.0 Australia licence</u>.

This document is valid for teaching and examining until 31 December 2013.

2011/38829(v3) Chemistry Data Sheet updated December 2012 Ref: 12-022

Periodic table

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 H hydrogen 1.008																	He helium 4.003
3	4											5	6	7	8	9	10
Li lithium	Be beryllium											B	C	N nitrogen	Oxygen	F fluorine	Ne
6.941	9.012											10.81	12.01	14.01	16.00	19.00	20.18
Na sodium	Mg magnesium											13 A{ aluminium	Si silicon	15 P phosphorus	16 S sulfur	17 Ce chlorine	18 Ar argon
22.99 19	24.31	21	20	22	l 24	25	26	1 07	T 20	20	1 20	26.98 31	28.09 32	30.97	32.06 34	35.45 35	39.95 36
K	Ca	Sc	22 Ti	23 V	Cr	Mn	Fe	Co	Ni Ni	Cu	Zn	Ğa	Ğe	As		Br	Kr
potassium 39.10	calcium 40.08	scandium 44.96	titanium 47.87	vanadium 50.94	chromium 52.00	manganese 54.94	iron 55.85	cobalt 58.93	nickel 58.69	copper 63.55	zinc 65.38	gallium 69.72	germanium 72.63	arsenic 74.92	Se selenium 78.96	bromine 79.90	krypton 83.80
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb rubidium 85.47	Sr strontium 87.62	yttrium 88.91	Zr zirconium 91.22	Nb niobium 92.91	Mo molybdenum 95.96	Tc technetium	Ru ruthenium 101.1	Rh rhodium 102.9	Pd palladium 106.4	Ag silver 107.9	Cd cadmium 112.4	In indium 114.8	Sn 118.7	Sb antimony 121.8	Te tellurium 127.6	iodine 126.9	Xe xenon 131.3
55	_56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs caesium 132.9	Ba barium 137.3	*La lanthanum 138.9	Hf hafnium 178.5	Ta tantalum 180.9	tungsten 183.8	Re rhenium 186.2	Os osmium 190.2	Ir iridium 192.2	Pt platinum 195.1	Au gold 197.0	Hg mercury 200.6	Te thallium 204.4	Pb lead 207.2	Bi bismuth 209.0	Po	At astatine	Rn
87	_88	89-103	104	105	106	107	108	109	110	111	112		114		_116		
Fr francium	Ra radium	**AC	Rf rutherfordium	Db dubnium	Sg seaborgium	Bh	Hs hassium	Mt meitnerium	DS darmstadtium	Rg roentgenium	Cn		Fe flerovium		LV		
												ı		1		ı	
Key:		* Lantl	hanida	58	59	60	61	62	63	64	65	66	67	68	69	70	71
		serie:	hanide s	Ce	Pr praseodymium	Nd neodymium	Pm	Sm samarium	Eu europium	Gd gadolinium	Tb terbium	Dy	Но	Er	Tm	Yb ytterbium	Lu lutetium
Atomic nu	ımber			140.1	140.9	144.2	promediani	150.4	152.0	157.3	158.9	dysprosium 162.5	holmium 164.9	167.3	168.9	173.1	175.0

Pu

95

Am

Cm

Bk

[Data source: The International Union of Pure and Applied Chemistry Periodic Table of the Elements (June 2012)]

U

uranium 238.0

Np

Colours of selected ionic substances

Es

Cf

In general, ionic solids have the same colour as that of any coloured ion they contain.

100

Fm

101

Md

102

No

103

Lr

Other coloured

substances

Two colourless ions in general produce a white solid. Selected exceptions to these two basic rules are noted below.

Ionic Solid	Colour
copper(II) carbonate	green
copper(II) chloride	green
copper(II) oxide	black
copper(II) sulfide	black
lead(II) iodide	yellow
lead(II) sulfide	grey
manganese(IV) oxide	black
silver carbonate	yellow
silver iodide	pale yellow
silver oxide	brown
silver sulfide	black

Most gases and liquids are colourless, and most metals are silvery or grey. Selected exceptions to these basic rules are noted below.

Substance	State	Colour
copper	solid	salmon pink
gold	solid	yellow
nitrogen dioxide	gas	brown
sulfur	solid	yellow

Coloured ions in aqueous solution

Cation	Colour
Cr ³⁺	deep green
Co ²⁺	pink
Cu ²⁺	blue
Fe ²⁺	pale green
Fe ³⁺	pale brown
Mn ²⁺	pale pink
Ni ²⁺	green

Anion	Colour
CrO ₄ ²⁻	yellow
Cr ₂ O ₇ ²⁻	orange
MnO ₄ -	purple

Number of moles

Number of moles of solute

Number of moles of a gas at STP n

PV = nRTIdeal gas law

ppm = $\frac{\text{mass of solute (mg)}}{\text{mass of solution (kg)}}$ Parts per million

pH of a solution $pH = -log[H^{+}]$

Units

Formulae

Volumes are given in the units of litres (L), or millilitres (mL)

Temperatures are given in the units of degrees Celsius (°C) or kelvin (K).

It may be assumed that 0.0 °C = 273.15 K

Energy changes are given in kilojoules (kJ)

Pressures are given in kilopascals (kPa)

Solution concentrations are given in the units moles per litre (mol L-1),

grams per litre (g L-1) or parts per million (ppm).

Constants

Universal gas constant, R = 8.314 J K⁻¹ mol⁻¹

Avogadro constant, N = 6.022×10^{23} mol⁻¹

Volume of 1.00 mol of an ideal gas at 0.0 °C and 100.0 kPa is 22.71 L

S.T.P. is 0.0 °C and 100.0 kPa

Equilibrium constant for water at 25 °C, $K_w = 1.00 \times 10^{-14}$

Coloured halogens

Halogen	Colour of free element
F ₂ (g)	yellow
Cl ₂ (g)	greenish-yellow
Br ₂ (ℓ)	red
$I_2(s)$	purple

Halogen	Colour of halogen in			
	aqueous solution			
Cl ₂ (aq)	pale yellow			
Br ₂ (aq)	orange			
I ₂ (aq)	brown			

Halogen	Colour of halogen in organic solvent
Br ₂	red
I ₂	purple

Solubility rules for ionic solids in water Soluble in water

** Actinide

Symbol

Standard

atomic weight

Soluble	Exceptions				
	Insoluble	Slightly soluble			
Most chlorides	AgCl	PbCl ₂			
Most bromides	AgBr	PbBr ₂			
Most iodides	AgI, PbI ₂				
All nitrates	No exceptions				
All ethanoates	NO CACCE	70013			
Most sulfates	SrSO ₄ , BaSO ₄ , PbSO ₄	CaSO ₄ , Ag ₂ SO ₄			

Th

Pa

orotactinium 231.0

Insoluble in water

Insoluble	Exceptions	
	Soluble	Slightly soluble
Most hydroxides	NaOH, KOH, Ba(OH) ₂ (note: NH ₄ OH and AgOH do not exist)	Ca(OH) ₂ , Sr(OH) ₂
Most carbonates	Na ₂ CO ₃ , K ₂ CO ₃ , (NH ₄) ₂ CO ₃	
Most phosphates	Na ₃ PO ₄ , K ₃ PO ₄ , (NH ₄) ₃ PO ₄	
Most sulfides	Na ₂ S, K ₂ S, (NH ₄) ₂ S	

Soluble = more than 0.1 mole dissolves per litre Slightly soluble = between 0.01 and 0.1 mole dissolves per litre Insoluble = less than 0.01 mole dissolves per litre