

K_W and pH CALCULATIONS

Define 'pH' as $-\log_{10}[\text{H}^+_{(\text{aq})}]$ and calculate the pH of strong acid solutions and strong base solutions.

Since the K_W for water = $[\text{H}^+_{(\text{aq})}] \times [\text{OH}^-_{(\text{aq})}] = 10^{-14}$ at 298 K (25^o C) it follows that

$$[\text{H}^+_{(\text{aq})}] = 10^{-14} \div [\text{OH}^-_{(\text{aq})}] \quad \text{or} \quad [\text{OH}^-_{(\text{aq})}] = 10^{-14} \div [\text{H}^+_{(\text{aq})}]$$

alternatively

$$\text{pH} + \text{pOH} = 14$$

$$\text{pH} = 14 - \text{pOH} \quad \text{or} \quad \text{pOH} = 14 - \text{pH}$$

Remember that an equilibrium constant is temperature dependant, therefore K_W will have a different value at temperatures other than 298 K (25^o C). Water will however be neutral as the $[\text{H}^+_{(\text{aq})}] = [\text{OH}^-_{(\text{aq})}]$ even if they are $\neq 10^{-7} \text{ mol L}^{-1}$.

$$K_W = [\text{H}^+] [\text{OH}^-] = 10^{-14} \text{ or } \text{pH} + \text{pOH} = 14$$

Set 1.

Unless stated otherwise the, the temperature is 298 K (25^o C)

- Q1. Calculate the concentration of OH^- ions in 0.10 mol L^{-1} HCl.
- Q2. Show that the concentration of H^+ (H_3O^+) ions in pure water is $1 \times 10^{-7} \text{ mol L}^{-1}$.
- Q3. Calculate the $[\text{H}^+]$ in a 0.25 mol L^{-1} sodium hydroxide.
- Q4. 3.65 grams of HCl gas are dissolved in enough water to make 1.5 L of solution. Calculate for this solution
- A. the concentration of the solution
 - B. $[\text{H}^+]$
 - C. $[\text{OH}^-]$
- Q5. A solution contains 11.22 grams of potassium hydroxide in 250 mL of solution. Calculate for this solution
- A. the concentration of the solution
 - B. $[\text{H}^+]$
 - C. $[\text{OH}^-]$
- Q6. For a 0.02 mol L^{-1} of nitric acid, calculate the $[\text{OH}^-]$ at
- A. 25 ^oC
 - B. 0 ^oC ($K_W = 1.1 \times 10^{-15}$)
- Q7. Explain why for pure water, acidic, basic and salt solutions the K_W for water at 25 ^oC is always 1.0×10^{-14} .
- Q8. Concentrated hydrochloric acid has a concentration of 11.7 mol L^{-1} . Calculate the pH and the $[\text{OH}^-]$ in this solution in mol L^{-1} .

Set 2.

Q1. Calculate the pH of each of the following solutions:

- | | | | |
|----|---|----|--|
| A. | $0.1 \text{ mol L}^{-1} \text{ HCl}$ | B. | $0.25 \text{ mol L}^{-1} \text{ HNO}_3$ |
| C. | $0.002 \text{ mol L}^{-1} \text{ Ba(OH)}_2$ | D. | $7.3 \text{ g L}^{-1} \text{ HCl}$ |
| E. | $6.3 \text{ g} / 250 \text{ mL HNO}_3$ | F. | $0.55 \text{ mol L}^{-1} \text{ HCl}$ |
| G. | $11.7 \text{ mol L}^{-1} \text{ HCl}$ | H. | $1.25 \times 10^{-5} \text{ mol L}^{-1} \text{ H}^+$ |

Q2. For a 0.10 mol L^{-1} solution of NaOH at $25 \text{ }^\circ\text{C}$ calculate the:

- A. $[\text{OH}^-]$
- B. $[\text{H}^+]$
- C. pH

Q3. 8.0 grams of NaOH is dissolved 5.0 L of solution at $25 \text{ }^\circ\text{C}$. Calculate the pH of this solution.

Q4. 0.561 grams of KOH is dissolved in 200 mL of solution. Calculate the pH

Q5. Calculate the pH of a $6.5 \times 10^{-4} \text{ mol L}^{-1} \text{ Ca(OH)}_2$ at $25 \text{ }^\circ\text{C}$.

Q6. The pH of vinegar is about 2.8 at $25 \text{ }^\circ\text{C}$. Calculate $[\text{H}^+]$.

Q7. The pH of human blood is about 7.4. Calculate $[\text{H}^+]$ and $[\text{OH}^-]$ (assume $25 \text{ }^\circ\text{C}$).

Q8. Calculate the $[\text{H}^+]$ and the $[\text{OH}^-]$ in a $0.3 \text{ mol L}^{-1} \text{ HCl}$ at $25 \text{ }^\circ\text{C}$.

Q9. A solution of KOH is made by dissolving 1.06×10^{-5} grams in 300 mL of solution. Calculate the pH of this solution at $25 \text{ }^\circ\text{C}$ and state whether the solution is slightly acidic or slightly basic (alkaline).

Q10. The average pH of sea-water at $25 \text{ }^\circ\text{C}$ is 8.5. Calculate the $[\text{H}^+]$ and the $[\text{OH}^-]$.

Q11. The pH of stomach acid is 1.7. Calculate the $[\text{H}^+]$ and the $[\text{OH}^-]$ in the stomach.

Set 3.

- Q1. What is the pH of each of the following solutions
- A. 0.01 mol L⁻¹ HCl
 - B. 0.1 mol L⁻¹ solution of a monoprotic acid which is 20% ionised.
 - C. A solution of HCl containing 2 g of HCl per litre.
 - D. A solution containing 2 g NaOH per litre.
 - E. A solution containing 0.63 g of HNO₃ in 500 mLs of solution.
 - F. A 0.01 mol L⁻¹ solution of ethanoic acid (CH₃COOH) given that it is 4.2% ionised at this concentration.
- Q2. Calculate the hydrogen ion concentration of solutions whose pH values are
- A. 4.3
 - B. 10.7
 - C. 7
 - D. 0
- Q3. A. Calculate the pH of a solution obtained by adding 49 mLs of 0.15 mol L⁻¹ NaOH to 50 mLs of 0.12 mol L⁻¹ HCl.
- B. Calculate the pH of a solution obtained by adding 19.4 mLs of 0.072 mol L⁻¹ Ba(OH)₂ to 27.8 mLs of 0.058 mol L⁻¹ HCl.
- Q4. Arrange the following 0.1 M solutions in order of increasing pH
- NaCl NH₄Cl NaOH HCl NH₃ CH₃COOH
- Q5. A. Explain why the pH of 0.1 mol L⁻¹ HCl is 1.0 while that of 0.1 mol L⁻¹ CH₃COOH is 2.87.
- B. Why is a solution of iron(III) chloride acidic?
- Q6. Calculate the pH of the solutions formed when 50 mLs of 0.1 mol L⁻¹ HCl is added to each of the following
- A. 49.5 mLs 0.23 mol L⁻¹ NaOH
 - B. 25.0 mLs 0.15 mol L⁻¹ Ba(OH)₂
 - C. 13.6 mLs 0.042 mol L⁻¹ KOH
 - D. 24.7 mLs 0.059 mol L⁻¹ HNO₃
 - E. 14.8 mLs 0.037 mol L⁻¹ NaOH and 15.0 mLs 0.14 mol L⁻¹ Ca(OH)₂

Answers:

Set 1

1. $1 \times 10^{-13} \text{ molL}^{-1}$
2. $[\text{H}^+][\text{OH}^-] = 10^{-14}$; $[\text{H}^+] = [\text{OH}^-]$; $[\text{H}^+]^2 = 10^{-14}$; $\sqrt{[\text{H}^+]^2} = [\text{H}^+] = \sqrt{10^{-14}} = 10^{-7}$
3. $4 \times 10^{-14} \text{ molL}^{-1}$
4. A. 0.067 molL^{-1} B. $1.5 \times 10^{-13} \text{ molL}^{-1}$
5. A. 0.8 molL^{-1} B. $1.25 \times 10^{-14} \text{ molL}^{-1}$
6. A. $5 \times 10^{-13} \text{ molL}^{-1}$ B. $5.5 \times 10^{-14} \text{ molL}^{-1}$
7. $K_w = [\text{H}^+][\text{OH}^-] = 10^{-14}$. The value of the equilibrium constant is constant (same) at a specified temperature.
8. $\text{pH} = -1.07$; $[\text{OH}^-] = 8.55 \times 10^{-16} \text{ molL}^{-1}$

Set 2

1. A. 1.0 B. 0.60 C. 11.6 D. 1.18
E. 0.40 F. 0.26 G. -1.07 H. 4.9
2. A. 0.1 B. 1×10^{-13} C. 11.6
3. A. 0.04 molL^{-1} B. $2.5 \times 10^{-13} \text{ molL}^{-1}$ C. 12.6
4. 12.7
5. 11.1
6. $1.58 \times 10^{-13} \text{ molL}^{-1}$
7. $[\text{H}^+] = 3.98 \times 10^{-8} \text{ molL}^{-1}$; $[\text{OH}^-] = 2.51 \times 10^{-7} \text{ molL}^{-1}$
8. $[\text{H}^+] = 0.3 \text{ molL}^{-1}$; $[\text{OH}^-] = 3.33 \times 10^{-14} \text{ molL}^{-1}$
9. 7.8; slightly alkaline
10. $[\text{H}^+] = 3.16 \times 10^{-8} \text{ molL}^{-1}$ $[\text{OH}^-] = 3.16 \times 10^{-8} \text{ molL}^{-1}$
11. $[\text{H}^+] = 0.02 \text{ molL}^{-1}$; $[\text{OH}^-] = 5.0 \times 10^{-13} \text{ molL}^{-1}$

Set 3

1. A. 2.00 B. 1.70 C. 1.26 D. 12.7 E. 1.70 F. 3.38
2. A. 5.01×10^{-5} B. 2.00×10^{-11}
C. 10^{-7} D. 10^0 or 1
3. A. 12.1 B. 12.4
4. HC/ CH_3COOH $\text{NH}_4\text{C/}$ NaC/ NH_3 NaOH
5. A. The $[\text{H}^+]$ in HC/ = 0.1 molL^{-1} because it is fully dissociated into ions whereas the $[\text{H}^+]$ in $\text{CH}_3\text{COOH} = 0.00132 \text{ molL}^{-1}$ because it is only partially dissociated into ions, much of the ethanoic acid remaining as molecular CH_3COOH .
B. Fe^{3+} ions react with water forming H^+ ions according to the following equation
 $\text{Fe}^{3+}_{(\text{aq})} + 3\text{H}_2\text{O}_{(\text{l})} \rightarrow \text{Fe}(\text{OH})_{3(\text{s})} + 3\text{H}^+_{(\text{aq})}$
6. A. 12.8 B. 12.5 C. 1.16 D. 1.06 E. 2.50

Set 1

$$K_w = 10^{-14} = [OH^-][H^+]$$

Q1 $10^{-14} = [OH^-][0.1]$
 $= 10^{-13} \text{ mol L}^{-1}$

Q2 $10^{-14} = [OH^-][H^+]$ pure water neutral
so $10^{-14} = [10^{-7}][10^{-7}]$

Q3 $10^{-14} = [0.25][H^+]$
 $= 4 \times 10^{-14}$

Q4 a) $n(\text{KCl}) = \frac{m}{M_r} = \frac{3.65}{36.45} = 0.1 \text{ mol}$

$$n(\text{KCl}) = CV$$
$$0.1 = C \times 1.5 \quad C_{(\text{KCl})} = 0.067 \text{ mol L}^{-1}$$

b) $C_{(\text{KCl})} = C_{(H^+)} = 0.067 \text{ mol L}^{-1}$

c) $10^{-14} = [H^+][OH^-]$
 $10^{-14} = [0.067][OH^-]$
 $[OH^-] = 1.5 \times 10^{-13} \text{ mol L}^{-1}$

Q5 a) $n(\text{KOH}) = \frac{m}{M_r} = \frac{11.22}{56.1} = 0.2 \text{ mol}$

$$n(\text{KOH}) = CV \quad 0.2 = C \times 0.25 = \underline{0.8 \text{ mol L}^{-1}}$$

c) $C_{(\text{KOH})} = C_{(OH^-)} = 0.8 \text{ mol L}^{-1}$

b) $10^{-14} = [H^+][OH^-]$
 $10^{-14} = [H^+][0.8]$
 $[H^+] = 1.25 \times 10^{-14} \text{ mol L}^{-1}$

Q6

a) $C(\text{HNO}_3) = C(\text{H}^+)$

$$10^{-14} = [\text{H}^+][\text{OH}^-]$$

$$10^{-14} = [0.02][\text{OH}^-]$$

$$[\text{OH}^-] = 5 \times 10^{-13} \text{ mol L}^{-1}$$

b) $1.1 \times 10^{-15} = [\text{H}^+][\text{OH}^-]$

$$= [0.02][\text{OH}^-]$$

$$[\text{OH}^-] = 5.5 \times 10^{-14} \text{ mol L}^{-1}$$

Q8

$$\text{pH} = -\log_{10} [11.7]$$

$$= -1.07$$

$$10^{-14} = [\text{H}^+][\text{OH}^-]$$

$$= [11.7][\text{OH}^-]$$

$$= 8.55 \times 10^{-16} \text{ mol L}^{-1}$$

Set 2

Q1

a) $pH = -\log [H^+] = -\log_{10} [0.1] = 1$

b) $-\log [0.25] = 0.6$

c) $0.002 \text{ mol of } Ba(OH)_2 = 0.004 \text{ mol of } OH^-$

$$\begin{aligned} \text{So } 10^{-14} &= [H^+][OH^-] \\ &= [H^+][0.004] \\ &= 2.5 \times 10^{-12} \cdot \log_{10} = 11.6 \end{aligned}$$

d) 7.3 g L^{-1} of HCl

$$\text{g L}^{-1} \div M \rightarrow \text{mol L}^{-1}$$

$$7.3 \div 36.45 = 0.2 \text{ mol L}^{-1}$$

$$pH = -\log [0.2]$$

$$pH = 0.7$$

e) $6.3 \text{ g} / 250 \text{ ml}$ so $\times \times = 25.2 \text{ g L}^{-1}$

$$\frac{25.2}{M_r (63)} = \text{mol L}^{-1} \quad \text{so} = 0.4 \text{ mol L}^{-1}$$

$$\begin{aligned} M_r &= H_2O_3 \\ &= 63 \end{aligned}$$

$$pH = -\log [0.4]$$

$$= 0.4$$

f) $pH = -\log [0.55] \quad pH = 0.26$

g) $pH = -\log [11.7] \quad pH = -1.07$

h) $pH = -\log [1.25 \times 10^{-5}] \quad pH = 4.9$

Q2 b.) $10^{-14} = [OH^-][H^+] = [0.1][H^+] = 1 \times 10^{-13} \text{ mol L}^{-1}$

a) 0.1

c) $pH = 13$.

Q3

$$n(\text{NaOH}) = 8 \text{ g per } 5 \text{ L so } 1.6 \text{ g L}^{-1}$$

$$1.6 \text{ g L}^{-1} \div 40 = 0.04 \text{ mol L}^{-1}$$

$$10^{-14} = [\text{OH}^-][\text{H}^+] = 10^{-14} = [0.04][\text{H}^+]$$

$$[\text{H}^+] = 2.5 \times 10^{-13} \text{ mol L}^{-1} \text{ pH} = 12.6$$

Q4

$$n(\text{KOH}) = 0.561 \text{ g per } 200 \text{ mL, so } 0.561 \times 5 = 2.805 \text{ g L}^{-1}$$

$$2.805 \text{ g L}^{-1} \div 56.1 = 0.05 \text{ mol L}^{-1}$$

$$10^{-14} = [0.05][\text{H}^+] \quad [\text{H}^+] = 2 \times 10^{-13} \text{ mol L}^{-1} \text{ pH} = 12.70$$

Q5

$$n(\text{Ca(OH)}_2) \times 2 = n(\text{OH}^-) \quad 6.5 \times 10^{-4} \times 2 = 0.0013 \text{ mol L}^{-1}$$

$$10^{-14} = [0.0013][\text{H}^+] = 0.0013 [\text{H}^+] \quad [\text{H}^+] = 7.69 \times 10^{-12} \text{ mol L}^{-1}$$

$$\text{pH} = 11.11$$

Q6

$$\text{pH} = 2.8 \quad \text{so} \quad 2.8 = -\log[\text{H}^+]$$

$$10^{-2.8} = 1.58 \times 10^{-3} \text{ mol L}^{-1}$$

Q7

$$\text{pH} = 7.4 \quad \text{so} \quad 10^{-7.4} = 3.98 \times 10^{-8} \text{ mol L}^{-1} = [\text{H}^+]$$

$$\text{so} \quad 10^{-14} = 3.98 \times 10^{-8} [\text{OH}^-] = 2.51 \times 10^{-7} \text{ mol L}^{-1}$$

Q8

$$10^{-14} = [0.3][\text{OH}^-] = 3.33 \times 10^{-14} \text{ mol L}^{-1}$$

Q9

$$1.06 \times 10^{-5} \times \frac{1000}{300} = 3.53 \times 10^{-5} \text{ g L}^{-1} \div 56.1 = 6.30 \times 10^{-7} \text{ mol L}^{-1}$$

$$10^{-14} = 6.3 \times 10^{-7} [\text{H}^+] = 1.587 \times 10^{-8} = \text{pH } 7.8 \text{ bain}$$

Q10

$$8.5 = -\log[\text{H}^+] = 3.16 \times 10^{-9} \text{ mol L}^{-1}, \text{ so } 10^{-14} = [\text{OH}^-] 3.16 \times 10^{-9} = 3.16 \times 10^{-6} \text{ mol L}^{-1}$$

Q11

$$11.7 = -\log[\text{H}^+] = 0.02 \text{ mol L}^{-1}, \text{ so } 10^{-14} = [\text{OH}^-] 0.02 = 5 \times 10^{-13} \text{ mol L}^{-1}$$

Set 3

Q1

- a) $pH = -\log(0.01) = 2$
- b) $0.2 \times 0.1 = 0.02 \text{ mol l}^{-1} = 1.70$
- c) $2 \text{ g l}^{-1} \div M_r \cdot 36.45 = 0.055 \text{ mol l}^{-1} = 1.26 \text{ pH}$
- d) $2 \text{ g l}^{-1} \div 40 = 0.05 \text{ mol l}^{-1}$, $10^{-14} = [H^+][0.05]$
 $= 2 \times 10^{-3} = \text{pH } 12.70$
- e) $0.63 \text{ g in } 0.5 \text{ L} = 1.26 \text{ g l}^{-1} \div 63 = 0.02 \text{ mol l}^{-1} = 1.7 \text{ pH}$
- f) $0.01 \times 4\% = 0.0004 \text{ mol l}^{-1} \text{ pH} = 3.38$

Q2

- a) $pH = -\log[H^+] = 5 \times 10^{-5} \text{ mol l}^{-1}$
- b) $10.7 \times -1 = -10.7 \Rightarrow 10^x = 2 \times 10^{-11} \text{ mol l}^{-1}$
- c) $1 \times 10^{-7} \text{ mol l}^{-1}$ d) 1 mol l^{-1}

Q3

- a)

<u>NaOH</u>	<u>KCl</u>
$n = cV$	$n = cV$
$= 0.115 \times 0.1049$	$= 0.12 \times 0.105$
$= 0.00735 \text{ mol}$	$= 0.006 \text{ mol}$

LR KCl, excess OH- ions $0.00735 - 0.006 = 0.00135 \text{ mol}$

$n(OH^-) = cV$ $10^{-14} = [H^+] 0.0136$
 $0.00135 = c \times 0.099$ $K^+ = 7.35 \times 10^{-13} \text{ mol l}^{-1} \text{ pH} = 12.1$
 $c(OH^-) = 0.0136 \text{ mol l}^{-1}$

b)

- | | |
|--|--|
| <u>Ba(OH)₂</u> | <u>KCl</u> |
| $n = cV$ | $n = cV$ |
| $n = 0.072 \times (19.4 \times 10^{-3})$ | $n = 0.058 \times (27.8 \times 10^{-3})$ |
| $n = 0.0013968 \text{ mol}$ | $n = 0.0016124$ |
| $n(OH^-) = 2 \times n(Ba(OH)_2)$ | \therefore KCl LR |
| $= 0.0027936 \text{ mol}$ | |
- excess OH- ions $0.0027936 - 0.0016124 = 0.0011812 \text{ mol}$
- $n(OH^-) = c \times 0.0472$ $10^{-14} = [H^+] 0.025$
 $= 0.025 \text{ mol l}^{-1} \rightarrow 4 \times 10^{-13} = 12.4 \text{ pH}$

Q6 B

$$\lambda(\text{HCl}) = 0.005 \text{ m} = \lambda(\text{H}^+)$$

$$\lambda(\text{HCl}) = cV = 0.1 \times (5 \times 10^{-3})$$

$$\lambda(\text{Ba(OH)}_2) = cV = 0.15 \times (25 \times 10^{-3}) = 0.00375 \text{ m}$$

$$\lambda(\text{Ba(OH)}_2) \times 2 = \lambda(\text{OH}^-)$$

$$= 0.0075 \text{ m}$$

$$\lambda(\text{OH}^-)_{\text{excess}} = 0.0075 - 0.005 = 0.0025$$

$$\lambda(\text{OH}^-) = 0.0025$$

$$\lambda(\text{OH}^-) = cV$$

$$= c \times 0.075$$

$$c[\text{OH}^-] = 0.0333 \text{ m}$$

$$10^{-14} = [\text{OH}^-][\text{H}^+]$$

$$\frac{10^{-14}}{0.0333} = 3 \times 10^{-13} \quad \text{pH} = 12.5$$

Q6 C

$$\lambda(\text{KOH}) = cV = 0.042 \times (3.6 \times 10^{-3}) = 0.0005712 \text{ m}$$

$$\lambda(\text{HCl}) - \lambda(\text{KOH}) = 0.005 - 0.0005712$$

$$\lambda(\text{HCl}) = 0.0044288 = \lambda(\text{H}^+)$$

$$\lambda(\text{H}^+) = cV$$

$$0.0044288 = c \times 0.0636$$

$$c[\text{H}^+] = 0.0696352 \quad \text{pH} = 1.16$$

Q6 d

$$\lambda(\text{HNO}_3) = cV = 0.054 \times (2.7 \times 10^{-3}) = 0.0014573 \text{ m}$$

$$\lambda(\text{HNO}_3) + \lambda(\text{HCl}) =$$

$$0.0014573 + 0.005 = 0.0064573$$

$$\lambda(\text{H}^+) = cV$$

$$0.0064573 = c \times 0.0747$$

$$c[\text{H}^+] = 0.0864431$$

$$\text{pH} = 1.06$$

$$\frac{1 \text{ F}}{96485} n(\text{HNO}_3) = \frac{m}{M_r} = \frac{0.63}{63} = 0.01 \text{ mol}$$

$$n(\text{HNO}_3) = n(\text{H}^+)$$

$$n(\text{H}^+) = cV$$

$$\frac{n(\text{H}^+)}{V} = c = \frac{0.01}{0.5} = 0.02 \text{ mol L}^{-1}$$

$$\text{pH} = 1.7$$

set 3 6 A

$$n(\text{NaOH}) = cV = 0.23 \times (49.5 \times 10^{-3})$$

$$n(\text{OH}^-) = 0.011385 \text{ mol}$$

$$n(\text{HCl}) = cV = 0.1 \times (50 \times 10^{-3})$$

$$n(\text{H}^+) = 0.005 \text{ mol}$$

$$n(\text{OH}^-)_{\text{excess}} = 0.011385 - 0.005$$

$$= 0.006385 \text{ mol}$$

$$n(\text{OH}^-) = cV$$

$$c(\text{OH}^-) = \frac{n}{V} = \frac{0.006385}{(99.5 \times 10^{-3})}$$

$$V_{\text{total}} = 50 \text{ ml} + 49.5 \text{ ml}$$

$$= 0.06471 \text{ mol L}^{-1}$$

$$[\text{OH}^-][\text{H}^+] = 10^{-14}$$

$$[\text{H}^+] = \frac{10^{-14}}{0.06471}$$

$$[\text{H}^+] = 1.5583 \times 10^{-13} \text{ M}$$

12.80734893

pH 12.8 . to 3 sf.

Q6 E

$$n(\text{NaOH}) = CV \\ = 0.1037 \times (14.8 \times 10^{-3})$$

$$n(\text{OH}^-) = 5.1476 \times 10^{-4} \text{ mol.}$$

$$n(\text{Ca(OH)}_2) = CV \\ = 0.14 \times (15 \times 10^{-3}) \\ = 0.0021 \text{ mol}$$

$$n(\text{Ca(OH)}_2) \times 2 = n(\text{OH}^-) \\ = 0.0042 \text{ mol.}$$

$$n(\text{OH}^-)_{\text{excess}} = 5.1476 \times 10^{-4} + 0.0042$$

$$= 4.7476 \times 10^{-3} \text{ mol.}$$

$$n(\text{H}^+) = 0.005 \text{ mol.} - 4.7476 \times 10^{-3}$$

$$n(\text{H}^+)_{\text{excess}} = 2.524 \times 10^{-4} \text{ mol}$$

$$n(\text{H}^+) = CV$$

$$n(\text{H}^+) = C$$

$$V = 79.8 \times 10^{-3}$$

$$C = 3.1629 \times 10^{-3}$$

2.4999 so $\text{pH} = 2.5$ to 2.5F.