How to do a Titration Calculation

Example: A 2.34g sample of an impure substance that is known to contain sodium hydroxide has been given to you. The sample was dissolved in distilled water and made up to the mark in a 250mL volumetric flask. 18.5mL of a 0.200M standardized solution of Hydrochloric acid, was required to react with 20mL aliquots of the sodium hydroxide solution. Determine the percentage purity of the sample.

Calculate:

(

- a. The number of moles of H⁺ which reacted.
- b. The number of moles of OH⁻ in the 20.0mL sample.
- c. The number of moles of OH⁻ in the 250.0mL solution.
- d. The mass of OH⁻ in the whole solution.
- e. The percentage purity of the sample.

Step 1 – Draw a diagram.



Step 2 – Determine the number of moles of H^{+} which reacted.

$$n(H^{+-}) = c \times V$$

= 0.200 x 0.0185

= <u>3.70 x 10⁻³ mol</u>

Step 3 – Use the equation to determine the number of moles of OH^{-} which reacted with the H^{+} . This is in the 20.0mL sample.

$$n(OH^{-})_{in \ 20mL} = 1/1 \times n(H^{+})$$
$$= 1 \times (3.70 \times 10^{-3})$$
$$= 3.70 \times 10^{-3} \text{ mol}$$

C)

Step 4 – We know the number of moles of OH^{-} in 20.0mL of sample. We want to know the number of moles of OH^{-} in the 250.0mL solution. We can use the concentration.

c(OH⁻) = n/V in 20mL = $3.70 \times 10^{-3} / 0.02$ = 0.185 mol.L^{-1} Then in 250.0mL n = c x V = $(0.185) \times 0.250$ = $4.62 \times 10^{-2} \text{ mol}$ We can combine these steps:

Since C_{in 250mL} = C_{in 20mL}

(

(

 $n_1/V_1 = n_2/V_2$

 $n(OH^{-})_{in 250mL} = 250/20 \times n(OH^{-})_{in 20mL}$

= 250/20 x 3.70 x 10⁻³

 $= 4.62 \times 10^{-2} mol$

Step 5 – We now have the number of moles of OH⁻ in the 250.0mL flask. We now determine the mass.

 $n(OH^{-}) = n(NaOH)$ $m(NaOH) = n \times M$ $M(n_{eolt}) = 40.1 \text{ g.mol}^{-1}$ $= (4.62 \times 10^{2}) \times 40.1$ = 1.85g

Step 6 – This is the mass of pure NaOH in the sample. We divide this by the sample mass to determine purity.

% purity = mass(pure) / mass(sample) x 100

= (1.85 / 2.34) x 100

= <u>79.2%</u>

 \bigcirc

Titrating NaOH and Acetic Acid

Procedure:

(

Step 1. Rinse a clean 250 mL conical (erlenmeyer) flask with water.

Step 1. Rinse a clean 25.00 mL pipette (pipet) with vinegar. Pipette 25.00 mL of vinegar into the 250 mL conical (erlenmeyer) flask.

Step 2. Add 2 drops of phenolphthalein indicator to the vinegar. (The solution will remain colourless)

Step 3. Rinse a clean 50.00 mL burette (buret) with standardised 1.00 mol L⁻¹ aqueous sodium hydroxide solution. Fill the burette (buret) with this standardised 1.00 mol L⁻¹ NaOH(aq).

Step 4. Set up the equipment as in the diagram on the right.

Step 5. Run NaOH(aq) from the burette (buret) into the conical (erlenmeyer) flask until the solution changes colour from colourless to pink.

Step 6. Repeat the titration carefully several times until concordant titres are achieved.

Sample Results

	Trial 1 / mL	Trial 2 / mL	Trial 3 / mL
Final volume of NaOH(aq)	21.82	21.79	21.81
Initial volume of NaOH(aq)	0.00	0.00	0.00
Titre /mL (volume of NaOH(aq) used)	21.82	21.79	21.81
Average Titre /mL	<u>21.82 + 21</u>	. <u>79 + 21.81</u> 3	= 21.81



Calculating the Concentration of Acetic Acid in Vingear in mol L⁻¹

a. Write the balanced chemical equation for the neutralisation reaction:

· •	word equation	acetic acid (ethanoic acid)	+ so	dium hydroxide	→ (sodium acetate sodium ethanoate)	+ \	water	
balance	ed chemical equation	CH₃COOH(aq)	+	NaOH(aq)	\rightarrow	CH₃COO⁻Na⁺(aq)	+	H₂O	

b. Extract all the relevant data from the experiment.

balanced chemical equation $CH_3COOH(aq) + NaOH(aq) \rightarrow CH_3COO Na^+(aq) + H_2O$

volume /mL	25.00	21.81
concentration /mol L ⁻¹	?	1.00

c. Check the data for consistency:

Concentrations are usually given in M or mol L⁻¹ but volumes are often given in mL. You will need to convert the mL to L for consistency. The easiest way to do this is to multiply the volume in mL x 10^{-3} (which is the same as dividing the volume in mL by 1000)

rolovant chacies	acid	base	
relevant species	CH₃COOH(aq)	NaOH(aq)	
volume /mL	25.00	21.81	
volume /L	25.00/1000 = 0.02500	21.81/1000 = 0.02181	
concentration /mol L ⁻¹	?	1.00	

 d. Calculate the moles of NaOH(aq), n(NaOH) moles = concentration in mol L⁻¹ x volume in L = n = c x V

> volume of NaOH(aq) = v(NaOH) = 0.02181 L concentration of NaOH(aq) = c(NaOH) = 1.00 mol L-1

moles NaOH(aq) = n(NaOH) = c(NaOH) x V(NaOH) n(NaOH) = 0.02181 x 1.00 = 0.02181 mol

e. Use the balanced chemical equation to determine the stoichiometric (mole) ratio of acid to base:

n(acid):n(base) that is n(CH₃COOH):n(NaOH) is 1:1

- f. Use the stoichiometric (mole) ratio to calculate the moles of acetic acid 1 mole of NaOH neutralises 1 mole of CH₃COOH therefore 0.02181 moles of NaOH neutralises 0.02181 moles of CH₃COOH moles of acetic acid = n(CH₃COOH) = 0.02181 mol
- g. From the volume of vinegar (acetic acid solution) and the moles of acetic acid, calculate its concentration (c) in mol L⁻¹ :

concentration (mol L⁻¹) = moles ÷ volume (L) concentration of acetic acid = moles of acetic acid ÷ volume of acetic acid in L

moles of acetic acid = $n(CH_3COOH) = 0.02181$ mol volume of acetic acid solution (vinegar) = $v(CH_3COOH) = 0.02500$ L concentration of acetic acid solution (vinegar) = $n(CH_3COOH) \div v(CH_3COOH)$ $c(CH_3COOH) = 0.02181 \div 0.02500 = 0.8724$ mol L⁻¹

1

(

Concentration of acetic acid in vinegar in mol L⁻¹ (molarity) is 0.8724 mol L⁻¹

