

## Science Department

## Year 12 Chemistry ATAR

# **Test 6: Organic Chemistry**

Name:

#### Instructions to Students:

- 1. One lesson permitted
- 2. Attempt all questions
- 3. Write in the spaces provided
- 4. Show all working when required
- 5. All answers to be in blue or black pen, diagrams in pencil.

Multiple Choice	Short Answer	TOTAL	Final Percentage
/10	/52	/62	

#### Section 1 – Multiple Choice

1. Which of the following is the correct IUPAC name for the structure shown?



- (a) *cis*-2,2-dimethylbut-2-ene
- (b) trans-2,3-dimethylbut-2-ene
- (c) hexene
- (d) 2,3-dimethylbut-2-ene
- 2. Which of the following is not a correctly named isomer of hexene?
  - (a) 2,3-dimethylbut-1-ene
  - (b) cyclohexane
  - (c) 1,3,4-trimethylprop-2-ene
  - (d) 2-methylpent-1-ene
- 3. What is the correct IUPAC name of the structure shown below?



- (a) 3-aminohexanal
- (b) 2-ethylpentanamide
- (c) 1-amino-2-ethylpentanal
- (d) 2-ethylpenan-1-amine
- 4. The structure below represents?



- (a) A primary alcohol
- (b) A secondary alcohol
- (c) A tertiary alcohol
- (d) A ketone

- 5. Which of the following alcohols could be used to produce butanoic acid?
  - (a) butan-2-ol
  - (b) 2-methylpropan-1-ol
  - (c) butan-1-ol
  - (d) butanone

6. Which of the following would not undergo an addition reaction?



7. The correct balancing coefficients for the equation below are:

 $\_ CH_3CHCHCH_2CH_3 + \_ O_2 \rightarrow \_ CO_2 + \_ H_2O$ 

- (a) 1, 8, 10, 5
- (b) 2, 15, 10, 10
- (c) 2, 16, 20, 10
- (d) 1, 7, 10, 10
- 8. Which of the following reaction would have products that could be identified by the use of universal indicator?
  - (a) butan-1-ol and acidified potassium permanganate
  - (b) hexan-1-ol and a limited amount of dilute acidified potassium dichromate
  - (c) butan-2-ol and acidified potassium permanganate
  - (d) 2-methylpentan-2-ol and and a limited amount of dilute acidified potassium dichromate

- 9. Which of the following families of organic compound is the least soluble in water?
  - (a) The esters
  - (b) The alcohols
  - (c) The carboxylic acids
  - (d) The primary amines
- 10. To form the ester pentyl propanoate you could react the following substances under appropriate conditions.
  - (a) pentan-1-ol and pentanoic acid
  - (b) propan-1-ol and pentanoic acid
  - (c) pentan-2-ol and propanal
  - (d) pentan-1-ol and propanoic acid

#### YOU MUST SHOW ALL HYDROGEN ATOMS IN YOUR STRUCTURAL DIAGRAMS

1. Give the IUPAC name of the following structures:



(6 marks)

2. Give the <u>full structural formula</u> for the following organic chemicals (include all Hydrogens – except for cyclic compounds):

(a) butan-2-one	(b) 4-ethyl-5-fluorohexan-2-one
(c) methyl butanoate	(d) ethyl 3-hydroxypentanoate
(e) cis-2,3-diiodo-4-methylhex-2-ene	(f) cyclohepta-1,4-diene
(g) 3,4,4,5-tetrachloro-2-methylpentan-2-ol	(h) 1,7-dibromocyloct-1-ene

(8 marks)

- 3. A student is asked to identify four organic liquids, contained in four separate flasks.
  - Octene
  - Hexan-3-ol
  - Hexan-3-one
  - Butanoic acid

The student has access to any chemicals and glassware required.

Describe the chemical tests that should be carried out, and the observations, that enable the liquids to be identified. (Note using an acid base indicator is not a test you may use)

### Include equations to justify the choice of tests.

(9 marks

- 4. DRAW and NAME the major **organic** PRODUCT or PRODUCTS in the following reactions assuming appropriate conditions. NB. No balancing is required.
  - (a) Pent-2-ene and hydrogen fluoride gas.

(b) Benzene and steam under appropriate reaction conditions.

(c) Ethanol and pentanoic acid with an acid catalyst.

(d) Hept-3-ene ignited in an oxygen rich atmosphere.

(e) But-2-ene is reacted **<u>completely</u>** with excess Bromine water.

5. A certain organic compound is known to contain only carbon, hydrogen and oxygen.

The compound was analysed as follows.

A 2.149 g sample was burned, and the carbon dioxide produced was bubbled through a barium hydroxide solution, producing 11.27 g of barium carbonate  $(BaCO_3)$ .

 $CO_2$  +  $Ba(OH)_2 \rightarrow BaCO_3$  +  $H_2O$ 

The mass of water produced by burning of the sample was 0.7721 g

The compound was found to have a molecular weight of 150.1 g mol<sup>-1</sup>

- a) What is the empirical formula of the compound? (10 marks)b) What is the molecular formula of the compound? (2 marks)
- c) The compound is also known to be a carboxylic acid; that is, containing one COOH group. Write the molecular formula in the form of  $C_XH_YO_ZCOOH$  (giving values for X, Y and Z).

(1 mark)

6. The following table shows the solubilities of two amines in water.

	Methyl amine	Dodecyl amine	
Amine	CH₃NH₂	$CH_3(CH_2)_{11}NH_2$	
Solubility (g/100 mL)	108	0.05	

Explain why their solubilities are so different. (Include a labelled diagram.)

(6 marks)


End of Test