



Christ Church
Grammar School

Year 12 Chemistry

Equilibrium Test 2020

Time allowed:

45 minutes

Name:

Teachers:

NMO

CEM

KLD

MLC


Mark =/47

SECTION 1

MULTIPLE CHOICE


10 marks

1. Which one of the options about the following reversible reaction is true?



- (a) $K_c = \frac{[\text{SO}_2]^2 [\text{O}_2]}{[\text{SO}_3]^2}$
- (b) K_c is constant under all reaction conditions.
- (c) Sulfur trioxide is being formed when the reaction is at equilibrium.
- (d) Adding a catalyst will increase the yield of sulfur trioxide.

Questions 2 and 3 refer to the following reaction:



2. The equilibrium expression, K_c , for the reaction above would be

- (a) $K_c = \frac{[\text{N}_2\text{O}_4]}{2[\text{NO}_2]}$
- (b) $K_c = \frac{2[\text{NO}_2]}{[\text{N}_2\text{O}_4]}$
- (c) $K_c = \frac{[\text{N}_2\text{O}_4]}{[\text{NO}_2]^2}$
- (d) $K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]}$

3. What will happen to the value of K_c in the reaction described in question 2, if the concentration of the N_2O_4 is doubled?

- (a) K_c would not be affected.
- (b) K_c would be halved.
- (c) K_c would be doubled.
- (d) K_c would increase by a factor of 4.


4. At constant temperature, the addition of a catalyst to an equilibrium system,

- (a) increases the concentration of the products at equilibrium.
- (b) increases the energy of the molecules so more can successfully collide.
- (c) lowers the amount of energy released in the overall reaction.
- (d) decreases the time required for equilibrium to be reached.

5. In which of the following systems will the mass of the products increase if the volume of the container is increased?

- (a) $2\text{NO}_2(\text{g}) \rightleftharpoons \text{N}_2\text{O}_4(\text{g})$
- (b) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$
- (c) $\text{H}_2\text{O}(\text{g}) + \text{C}(\text{s}) \rightleftharpoons \text{H}_2(\text{g}) + \text{CO}(\text{g})$
- (d) $\text{H}_2(\text{g}) + \text{F}_2(\text{g}) \rightleftharpoons 2\text{HF}(\text{g})$


6. Consider the following reaction at equilibrium:



Which one of the following changes will increase the concentration of $\text{NO}_2(\text{g})$ in the mixture when equilibrium is re-established?

- (a) decreasing the concentration of NO at constant temperature and pressure
- (b) decreasing the concentration of O_2 at constant temperature and pressure
- (c) increasing the volume
- (d) decreasing the temperature

Questions 7 and 8 refer to the following reaction:



The following changes can be made to the reaction:

- (I) Increase the concentration of $\text{NH}_3(\text{g})$
- (II) Increase the concentration of $\text{H}_2\text{O}(\text{g})$
- (III) Decrease the temperature


7. Which of the changes will increase the yield of products?

- (a) I only
- (b) I and II
- (c) I and III
- (d) I, II and III

8. Which of the following will increase the rate of the forward reaction when equilibrium is re-established?

- (a) II only
- (b) I and II
- (c) I only
- (d) I, II and III





9. Consider the following system initially at equilibrium.




The system is initially at equilibrium. At time t_1 , the temperature of the system was increased. Which of the following best represents the changes in the forward and reverse reaction rates until equilibrium is re-established at time, t_2 ?

The forward reaction rate is represented by _____

The reverse reaction rate is represented by _____



10. Lead iodide is slightly soluble in water and the following equilibrium is produced:



Some $\text{Na}_2\text{SO}_4(\text{aq})$ was then added to the system and allowed to reach equilibrium. What would you expect to observe during this period?

- (a) No visible change.
- (b) White solid forms.
- (c) Yellow solid forms.
- (d) White solid forms and yellow solid decreases in mass.

SECTION 2


SHORT ANSWERS

37 marks

Question 11

21 marks

When potassium thiocyanate (KSCN) is mixed with iron(III) nitrate ($\text{Fe}(\text{NO}_3)_3$) in solution, an equilibrium mixture of Fe^{3+} , SCN^- , and the ion FeSCN^{2+} is formed according to the equation below. The formation of FeSCN^{2+} is exothermic.



- (a) Complete the table below using the terms *increase*, *decrease* or *no change* to indicate the effect of making the following changes to the system once equilibrium has been re-established. (12 marks)

| Change made | Rate of reverse reaction | Effect on $[\text{Fe}^{3+}]$ | Effect on K |
|--------------------------|--------------------------|------------------------------|-------------|
| Add FeCl_3 (s) | | | |
| Add H_2O | | | |
| Add a catalyst | | | |
| Decrease temperature | | | |


Question 12

11 marks

In the first stage of the production of nitric acid, ammonia is reacted with oxygen to produce nitrogen monoxide.


A hot platinum catalyst is used. The reaction is highly exothermic, and the heat given out by the reaction is sufficient to maintain the temperature needed to give a fast rate.

The system is at equilibrium until t_1 and then at t_1 , the volume of the vessel is reduced at constant temperature. Equilibrium is re-established at t_2 .



- (a) Sketch the rate-time graph for the forward and reverse reactions to show the changes in rate until the return to equilibrium.


Use forward ————— reverse ----- in your sketch below. (3 marks)



- (b) Using 'increase, decrease or no change' in the table below, identify the effect of reducing the volume on the mass and concentration of NH_3 and NO once equilibrium has been re-established. (4 marks)

| | NH_3 | NO |
|---------------|---------------|-------------|
| Mass | | |
| Concentration | | |

- (c) The second stage involves the reaction of nitrogen monoxide with more oxygen to form nitrogen dioxide.



At t_0 , 2.5 mol of NO and 1 mol of O_2 were injected into a 500 mL container.

At t_1 , equilibrium was reached with only 1 mol of NO remaining.

Use this information to construct a detailed graph for how the concentrations of NO, O_2 and NO_2 varied between t_0 and t_1 .

(4 marks)

