

MATHEMATICS METHODS

MAWA Semester 2 (Units 3 and 4) Examination 2016

Calculator-free

Marking Key

© MAWA, 2016

Licence Agreement

This examination is Copyright but may be freely used within the school that purchases this licence.

- The items that are contained in this examination are to be used solely in the school for which they are purchased.
- They are not to be shared in any manner with a school which has not purchased their own licence.
- The items and the solutions/markings keys are to be kept confidentially and not copied or made available to anyone who is not a teacher at the school. Teachers may give feedback to students in the form of showing them how the work is marked but students are not to retain a copy of the paper or marking guide until the agreed release date stipulated in the purchasing agreement/licence.

The release date for this exam and marking scheme is

- **the end of week 1 of term 4, 2016**

Section One: Calculator-free

(54 Marks)

Question 1(a)

Solution $\ln m = \frac{3}{2} \Rightarrow m = e^{\frac{3}{2}}$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> identifies correct base determines correct power 	1 1

Question 1(b)

Solution $\log[(m+3)m] = 1$ $(m+3)m = 10^1$ $m^2 + 3m - 10 = 0$ $(m+5)(m-2) = 0$ $m = -5$ or 2 but since m has to be greater than zero, $m = 2$ is the only solution.	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> applies logarithmic rule for a product correctly recognises base 10 creates equation with correct trinomial solves equation correctly giving the correct value of m 	1 1 1 1

Question 2(a) (i)

Solution $\frac{dy}{dx} = \frac{(6x^4 - x^3 + e)(4e^x) - (4e^x)(24x^3 - 3x^2)}{(6x^4 - x^3 + e)^2}$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> differentiates the 1st term on numerator correctly differentiates the 2nd term on numerator correctly squares factor on denominator 	1 1 1

Question 2(a)(ii)

<p>Solution</p> $\frac{dy}{dx} = \frac{d}{dx} [\ln(5x^3 + 3) - \ln(\sin(x))]$ $= \frac{d}{dx} [\ln(5x^3 + 3)] - \frac{d}{dx} [\ln(\sin(x))]$ $= \frac{15x^2}{(5x^3 + 3)} - \frac{\cos(x)}{\sin(x)}$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> • applies correctly logarithmic rule for quotients • differentiates correctly 1st term • differentiates correctly 2nd term 	<p>1</p> <p>1</p> <p>1</p>

Question 2(b)

<p>Solution</p> <p>Let $u = x^2 - \cos(x) \Rightarrow \frac{du}{dx} = 2x + \sin(x)$ and $\frac{dy}{du} = \frac{e^u}{2}$</p> $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} = \frac{e^u}{2} \times (2x + \sin(x)) = \frac{e^{x^2 - \cos(x)}}{2} (2x + \sin(x))$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> • differentiates correctly to determine 1st factor in chain rule • differentiates correctly to determine 2nd factor in chain rule • expresses $\frac{dy}{dx}$ in terms of x 	<p>1</p> <p>1</p> <p>1</p>

Question 3(a)

Solution Discrete random variable	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none">determines correct category	1

Question 3(b)

Solution Non-random variable	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none">determines correct category	1

Question 3(c)

Solution Continuous random variable	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none">determines correct category	1

Question 4

Solution $k \int_0^1 x - \frac{x^3}{3} dx = 1$ $k \left[\frac{x^2}{2} - \frac{x^4}{12} \right]_0^1 = 1$ $k \left[\frac{1}{2} - \frac{1}{12} \right] = 1 \Rightarrow k = \frac{12}{5}$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> • sets up integral and equates to one • integrates correctly • evaluates integral correctly • calculates the value of k 	<p>1</p> <p>1</p> <p>1</p> <p>1</p>

Question 5

Solution $p(1-p) = \left(\frac{\sqrt{3}}{4} \right)^2 = \frac{3}{16}$ $16p^2 - 16p + 3 = 0$ $(4p-1)(4p-3) = 0 \Rightarrow p = \frac{1}{4} \text{ or } p = \frac{3}{4}$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> • sets up equation using variance of a Bernoulli distribution • derives quadratic equation • factorises trinomial • solves correctly for p 	<p>1</p> <p>1</p> <p>1</p> <p>1</p>

Question 6(a)

Solution	
Function is valid for $x > -3$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> correctly states the values of x for which the function is valid 	1

Question 6(b)

Solution	
$\frac{dy}{dx} = \frac{2}{2x+6} = 4 \Rightarrow \frac{2x+6}{2} = \frac{1}{4} \Rightarrow x+3 = \frac{1}{4} \Rightarrow x = -2.75$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> differentiates correctly solves equation correctly 	1 1

Question 7(a)

Solution						
y	0	1	2	3	4	
$P(Y = y)$	0	k	$4k$	$9k$	$16k$	
Marking key/mathematical behaviours						Marks
<ul style="list-style-type: none"> correctly completes two values 						1
<ul style="list-style-type: none"> correctly completes 4 values 						1

Question 7(b)

Solution	
$k + 4k + 9k + 16k = 1$	
$30k = 1 \Rightarrow k = \frac{1}{30}$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> sums probabilities equal to one 	1
<ul style="list-style-type: none"> correctly solves equation for k 	1

Question 8

<p>Solution</p> $f(x) = \int f'(x) dx$ $= \int 2xe^{3x^2-1} dx$ $= \frac{1}{3}e^{3x^2-1} + c$ <p>since $f(0) = 0$:</p> $0 = \frac{1}{3}e^{-1} + c$ $c = -\frac{1}{3e}$ $f(x) = \frac{1}{3}e^{3x^2-1} - \frac{1}{3e}$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> determines indefinite integral 	1
<ul style="list-style-type: none"> substitutes initial conditions to calculate the constant c 	1
<ul style="list-style-type: none"> states $f(x)$ 	1

Question 9 (a)(i)

<p>Solution</p> $\hat{p} = \frac{20}{100} = \frac{1}{5}$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> determines the proportion 	1

Question 9(a)(ii)

Solution $E = 2 \times \sqrt{\frac{\frac{1}{5}(1-\frac{1}{5})}{100}}$ $= 2 \times \sqrt{\frac{4}{2500}}$ $= 2 \times \frac{2}{50}$ $= 0.08$ 95% CI is (0.12,0.28)	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> substitutes values for z, n and p simplifies square root simplifies E states interval 	1 1 1 1

Question 9(b)

Solution $E = 1 \times \sqrt{\frac{m(1-m)}{n_1}}$ 68% CI is $(m - \sqrt{\frac{m(1-m)}{n_1}}, m + \sqrt{\frac{m(1-m)}{n_1}})$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> determines E states confidence interval. 	1 1

Question 9(c)(i)


Solution	
n_2 is larger than n_1	
To increase confidence a larger interval is required for a stable sample size. Increasing n reduces the standard error and thus the interval can remain the same.	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> states n_2 is larger with reason states correct reason 	1
	1

Question 9(c)(ii)

Solution	
$E_1 = 1 \times \sqrt{\frac{m(1-m)}{n_1}}$ $E_2 = 1.5 \times \sqrt{\frac{m(1-m)}{n_2}}$	
Same interval so $E_1 = E_2$	
$\sqrt{\frac{m(1-m)}{n_1}} = 1.5 \times \sqrt{\frac{m(1-m)}{n_2}}$ $\frac{m(1-m)}{n_1} = (1.5)^2 \frac{m(1-m)}{n_2}$ $\frac{n_2}{n_1} = 2.25$ $n_2 = 2.25n_1$	
Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> equates E_1 and E_2 squares both sides states relationship 	1
	1
	1

Question 10(a)

Solution



$$\int_0^5 f(x)dx = \text{area of triangle (Area 1)}$$


$$= \frac{1}{2} \times 5 \times 9$$

$$= 22\frac{1}{2}$$

Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> identifies integral as area of correct triangle 	1
<ul style="list-style-type: none"> determines integral. 	1

Question 10(b)

Solution



$$\begin{aligned} \text{Area 2} &= \frac{1}{2} \times 2 \times 9 \\ &= 9 \text{ square units} \end{aligned}$$

$$\begin{aligned} \text{Area 3} &= 50 - 22\frac{1}{2} - 9 \\ &= 18\frac{1}{2} \text{ square units} \end{aligned}$$

$$\begin{aligned} \int_7^{10} f(x) dx &= - \text{Area 3} \\ &= -18\frac{1}{2} \end{aligned}$$

Marking key/mathematical behaviours	Marks
<ul style="list-style-type: none"> • calculates area 2 • calculates area 3 • determines integral 	<p>1</p> <p>1</p> <p>1</p>