




**Calculator Free
Applications of Differentiation**

Time: 45 minutes
Total Marks: 45
Your Score: / 45

Question One: [8 marks]

Sketch the graph of $f(x) = x^4 - 4x^2$ over the domain $-2 \leq x \leq 3$. Use calculus methods to determine the location and nature of any stationary points.



Mathematics Methods Unit 3

- (d) Determine the displacement of the particle when it changes direction for the second time. Comment on this result.
- (e) Determine when the particle reaches maximum velocity and calculate the speed of the particle at this time.

Question Three: [1, 1, 1, 2, 1, 1 = 7 marks]

A radioactive substance decays continuously at a rate of 2%. The amount of radioactive material remaining can be modelled by the function $A = A_0 e^{kt}$, where A is the amount of the substance in micrograms and t is the time in years.

- (a) State the value of k in this model.
- (b) Initially there are 20 micrograms of this substance. State the value of A_0 .
- (c) How many micrograms of the substance are there after 20 years?
- (d) Give an expression for the average rate of change of the amount of radioactive material in the first 10 years.
- (e) Determine an expression for the instantaneous rate of change of the amount of the radioactive substance.
- (f) Calculate the instantaneous rate of change of the amount of substance after 100 years.

Question Four: [1, 5 = 6 marks]

Consider the function $f(x) = \sqrt{x+3}$

- (a) Calculate $f(6)$.
- (b) Using your answer to (a) and the function $f(x)$, calculate the approximate value of $\sqrt{9.1}$

Question Five: [2, 2, 2, 2, 3 = 11 marks]

The current in a simple alternating current circuit is modelled by the function

$f(t) = 200\sin\left(\frac{\pi t}{60}\right)$, where t is the time in seconds and f is the flow of current in amps.

- (a) Calculate the amps after 10 seconds.

- (b) State the maximum amp flow and the time(s) at which these occur in the first minute.

- (c) Determine an expression for the instantaneous rate of change of the current.

- (d) Calculate the instantaneous rate of change of the current at 30 seconds.

- (e) Approximate the change in the flow of current in the 41st second.



SOLUTIONS
Calculator Free
Applications of Differentiation

Time: 45 minutes
 Total Marks: 45
 Your Score: / 45

Question One: [8 marks]

Sketch the graph of $f(x) = x^4 - 4x^2$ over the domain $-2 \leq x \leq 3$. Use calculus methods to determine the location and nature of any stationary points.

$$f'(x) = 4x^3 - 8x \quad \checkmark$$

$$4x^3 - 8x = 0$$

$$4x(x^2 - 2) = 0$$

$$x = 0, \sqrt{2}, -\sqrt{2} \quad \checkmark$$

$$f(0) = 0$$

$$f(\sqrt{2}) = 4 - 8 = -4 \quad \checkmark$$

$$f(-\sqrt{2}) = 4 - 8 = -4$$

$$f''(x) = 12x^2 - 8 \quad \checkmark$$

$$f''(0) < 0$$

$$\therefore (0, 0) \text{ max}$$

$$f''(\sqrt{2}) > 0$$


$$\therefore (\sqrt{2}, -4) \text{ min} \quad \checkmark$$

$$f''(-\sqrt{2}) > 0$$

$$\therefore (-\sqrt{2}, -4) \text{ min}$$

$$f(-2) = 0$$

$$f(3) = 45$$



Question Two: [1, 4, 3, 2, 3 = 13 marks]

The displacement of a particle moving in rectilinear motion is modelled by $x(t) = t(t-4)^2$, where t is the time in seconds and x is the displacement in metres.

- (a) Determine the initial displacement of the particle.

$$x(0) = 0m$$

- (b) Calculate the initial velocity of the particle and hence comment on the initial direction of the particle.

$$v(t) = (t-4)^2 + 2t(t-4)$$

$$v(0) = 16 + 0 = 16m/s$$

Therefore the particle is initially directed towards the right of the origin.

- (c) Determine when the particle first changes direction.

$$v(t) = 0$$

$$(t-4)(t-4+2t) = 0$$

$$(t-4)(3t-4) = 0$$

$$t = \frac{4}{3}, t = 4$$

$$\therefore t = \frac{4}{3}$$

Mathematics Methods Unit 3

- (d) Determine the displacement of the particle when it changes direction for the second time. Comment on this result.

$$x(4) = 4(4 - 4)^2 = 0m \quad \checkmark$$

When the particle changes direction for the second time it is located at the origin.



- (e) Determine when the particle reaches maximum velocity and calculate the speed of the particle at this time.

$$a(t) = 3(t - 4) + (3t - 4) \quad \checkmark$$

$$a(t) = 6t - 16$$

$$6t - 16 = 0$$

$$t = \frac{16}{6} = \frac{8}{3} \quad \checkmark$$

$$\left| v\left(\frac{8}{3}\right) \right| = \left| \left(\frac{8}{3} - 4\right)(8 - 4) \right| = 5\frac{1}{3} m/s \quad \checkmark$$

Question Three: [1, 1, 1, 2, 1, 1 = 7 marks]

A radioactive substance decays continuously at a rate of 2%. The amount of radioactive material remaining can be modelled by the function $A = A_0e^{kt}$, where A is the amount of the substance in micrograms and t is the time in years.

- (a) State the value of k in this model.

$$k = -0.02 \quad \checkmark$$

- (b) Initially there are 20 micrograms of this substance. State the value of A_0 .

$$A_0 = 20 \quad \checkmark$$

- (c) How many grams of the substance are there after 20 years?

$$A = 20e^{-0.02(20)} = 20e^{-0.4} \mu\text{g} \quad \checkmark$$

- (d) Give an expression for the average rate of change of the amount of radioactive material in the first 10 years.

$$= \frac{20e^{-0.2} - 20}{10} \quad \checkmark \checkmark$$

- (e) Determine an expression for the instantaneous rate of change of the amount of the radioactive substance.

$$\frac{dA}{dt} = -0.4e^{-0.02t} \quad \checkmark$$

- (f) Calculate the instantaneous rate of change of the amount of substance after 100 years.

$$\frac{dA}{dt} = -0.4e^{-0.02(100)} = -0.4e^{-2} \mu\text{g} / \text{year} \quad \checkmark$$

Question Four: [1, 5 = 6 marks]

Consider the function $f(x) = \sqrt{x+3}$

(a) Calculate $f(6)$.

$$f(6) = 3 \quad \checkmark$$

(b) Using your answer to (a) and the function $f(x)$, calculate the approximate value of $\sqrt{9.1}$

$$\partial x = 0.1 \quad \checkmark$$

$$f'(x) = 0.5(x+3)^{-0.5} \quad \checkmark$$

$$\partial y = f'(x)\partial x$$

$$\partial y = f'(6) \times 0.1 \quad \checkmark$$

$$\partial y = 0.5(9)^{-0.5} \times 0.1$$

$$\partial y = \frac{1}{60} \quad \checkmark$$

$$\therefore \sqrt{9.1} \approx 3\frac{1}{60} \quad \checkmark$$

Question Five: [2, 2, 2, 2, 3 = 11 marks]

The current in a simple alternating current circuit is modelled by the function

$f(t) = 200\sin\left(\frac{\pi t}{60}\right)$, where t is the time in seconds and f is the flow of current in amps.

- (a) Calculate the amps after 10 seconds.

$$f(10) = 100 \text{ amps} \quad \checkmark \checkmark$$

- (b) State the maximum amp flow and the time(s) at which these occur in the first minute.

$$\text{Maximum amp flow} = 200 \text{ amps} \quad \checkmark$$

$$t = 30, 90 \quad \checkmark$$

- (c) Determine an expression for the instantaneous rate of change of the current.

$$f'(t) = \frac{10\pi}{3} \cos\left(\frac{\pi t}{60}\right)$$

\checkmark
 \checkmark

- (d) Calculate the instantaneous rate of change of the current at 30 seconds.

$$f'(30) = \frac{10\pi}{3} \cos\left(\frac{30\pi}{60}\right) = 0 \text{ amps / sec} \quad \checkmark$$

\checkmark

- (e) Approximate the change in the flow of current in the 41st second.

$$f'(40) = \frac{10\pi}{3} \cos\left(\frac{40\pi}{60}\right) \quad \checkmark$$

$$= \frac{10\pi}{3} \times \frac{-1}{2}$$

$$= \frac{-5\pi}{3} \quad \checkmark$$