



Mercedes College

YEAR 12 MATHEMATICS METHODS Test 2 2016

Exponential and Trigonometric Functions

NAME: _____

Date: Tuesday 10th May

TEACHER: _____

Non-calculator section:	33 minutes	33 marks
Calculator section:	17 minutes	17 marks
OVERALL:	50 minutes	50 marks

INSTRUCTIONS:

Show FULL working Answer all questions on this test paper

Questions or parts of questions worth more than two marks require working to be shown to receive full marks.

Allowed: Maths Methods WACE formula sheets

TRIG FORMULA: $\cos 2\theta = \cos^2 \theta - \sin^2 \theta$ $\sin 2\theta = 2 \sin \theta \cos \theta$

$$\lim_{h \rightarrow 0} \left(\frac{1 - \cos h}{h} \right) = 0$$

$$\lim_{h \rightarrow 0} \left(\frac{\sin h}{h} \right) = 1$$

Q1 (5 marks)

Determine the equation of the tangent to the curve $y = \frac{\sin x}{x}$ at the point $(\pi, 0)$.

$$y = \frac{\sin x}{x}$$

$$u = \sin x \quad v = x$$

$$u' = \cos x \quad v' = 1$$

$$\frac{dy}{dx} = \frac{x \cos x - \sin x}{x^2}$$

$$\left. \frac{dy}{dx} \right|_{x=\pi} = \frac{\pi \cos \pi - \sin \pi}{\pi^2}$$

$$= \frac{\pi(-1) - 0}{\pi^2}$$

$$= -\frac{\pi}{\pi^2}$$

$$\frac{dy}{dx} = -\frac{1}{\pi}$$

LINEAR

$$y = mx + c$$

$$y = -\frac{1}{\pi}x + c$$

subst $(\pi, 0)$

$$0 = -\frac{1}{\pi}(\pi) + c$$

$$c = 1 \quad \checkmark$$

$$\boxed{y = -\frac{1}{\pi}x + 1}$$

Q2 (3 + 3 + 3 + 3 = 12 marks)

Determine $\frac{dy}{dx}$ for each of the following simplifying answers where possible.

(a) $y = e^{x^2-1} + 2 \cos(2x-1) + e^3$

$$\frac{dy}{dx} = 2x e^{x^2-1} - 4 \sin(2x-1)$$

(b) $y = \sin^3 5x = (\sin 5x)^3$

$$\begin{aligned} \frac{dy}{dx} &= 3 (\sin 5x)^2 (\cos 5x) 5 \\ &= 15 \cos 5x \sin^2 5x \end{aligned}$$

(c) $y = \frac{\cos x}{e^x}$

$$\begin{aligned} u &= \cos x & v &= e^x \\ u' &= -\sin x & v' &= e^x \end{aligned}$$

$$\frac{dy}{dx} = \frac{-\sin x e^x - \cos x e^x}{e^{2x}}$$

$$= \frac{-e^x (\sin x + \cos x)}{e^{2x}}$$

$$= -\frac{(\sin x + \cos x)}{e^x}$$

(d) $y = e^{(1-x)} \sin 2x$

$$u = e^{1-x} \quad v = \sin 2x$$

$$u' = -e^{1-x} \quad v' = 2 \cos 2x$$

$$\frac{dy}{dx} = -e^{1-x} \sin 2x + 2 \cos 2x e^{1-x}$$

$$= e^{1-x} (2 \cos 2x - \sin 2x)$$

Q3 (4 + 2 + 1 = 7 marks)
Evaluate the following.

(a) $\int_0^{\frac{\pi}{3}} (\cos \frac{x}{2} - \sin x) \cdot dx$

$$\begin{aligned} &= \left[2 \sin \frac{x}{2} + \cos 2x \right]_0^{\frac{\pi}{3}} \\ &= 2 \sin \frac{\pi}{6} + \cos \frac{\pi}{3} - 2 \sin 0 - \cos 0 \\ &= 2 \left(\frac{1}{2} \right) + \frac{1}{2} - 0 - 1 \\ &= \frac{1}{2} \end{aligned}$$

(b) $\frac{d}{dx} \left(\int_3^{x^2} e^{(\sqrt{t}-1)} \cdot dt \right)$ when $x = 2$

$$\begin{aligned} &= e^{\sqrt{x^2}-1} (2x) \\ &= e^{x-1} (2x) \\ \text{at } x=2 &= e^1 (4) \\ &= 4e \end{aligned}$$

(c) $\lim_{h \rightarrow 0} \frac{\sinh}{2h}$

$$\begin{aligned} &= \frac{1}{2} \lim_{h \rightarrow 0} \frac{\sinh}{h} \\ &= \frac{1}{2} (1) \\ &= \frac{1}{2} \end{aligned}$$

Q4 (2 + 2 + 3 + 2 = 9 marks)

Evaluate the following integrals.

(a) $\int (e^x + \cos x + \sin x) \cdot dx$

$$= e^x + \sin x - \cos x + C \quad \checkmark \checkmark$$

(b) $\int \frac{2}{e^{3x}} \cdot dx$

$$= \int 2e^{-3x} dx$$

$$= -\frac{2e^{-3x}}{3} + C \quad \checkmark \checkmark$$

(c) $\int \frac{\sin x \cos^3 x}{2} \cdot dx$

$$= \left(\frac{1}{-4}\right) \frac{1}{2} \int \boxed{-4 \sin x \cos^3 x dx}$$

$$= -\frac{1}{8} \cos^4 x + C$$

$\checkmark \checkmark \checkmark$

$$y = (\cos x)^4$$

$$\frac{dy}{dx} = -4 (\cos x)^3 \sin x$$

$$= -4 \sin x \cos^3 x$$

(d) $\int 4 \sin x \cos x \cdot dx$

$$= 2 \int 2 \sin x \cos x \cdot dx$$

$$= 2 \int \sin(2x) \cdot dx$$

$$= 2 \frac{(-\cos 2x)}{2} + C$$

$$= -\cos 2x + C \quad \checkmark \checkmark$$

END OF SECTION 1



Mercedes College

SOLUTIONS

YEAR 12 MATHEMATICS METHODS Test 2 2016

Exponential and Trigonometric Functions

NAME: _____

Date: Tuesday 10th May

TEACHER: _____

Non-calculator section:	33 minutes	33 marks
Calculator section:	17 minutes	17 marks
OVERALL:	50 minutes	50 marks

INSTRUCTIONS:

Show FULL working Answer all questions on this test paper

Questions or parts of questions worth more than two marks require working to be shown to receive full marks.

Allowed: Maths Methods WACE formula sheets, 3 calculators, 1 A4 page of notes

Q5 (4 marks)

A curve passes through the point $(\frac{\pi}{2}, \pi - 2)$ and has a gradient function given by

$\frac{dy}{dx} = 1 - 2 \cos x$. Determine the equation of the original curve.

$$y = x - 2 \sin x + c \quad \checkmark$$

Sub

$$\left(\frac{\pi}{2}, \pi - 2\right) \quad \pi - 2 = \frac{\pi}{2} - 2 \sin\left(\frac{\pi}{2}\right) + c \quad \checkmark$$

$$\pi - 2 = \frac{\pi}{2} - 2(1) + c$$

$$\frac{\pi}{2} = c \quad \checkmark$$

$$y = x - 2 \sin x + \frac{\pi}{2} \quad \checkmark$$

Q6 (1 + 2 + 1 + 2 + 4 = 10 marks)

The mass of a drug remaining in the bloodstream of a patient is changing according to the rule $\frac{dM}{dt} = -0.12M$, where M is the mass of drug remaining t hours after the initial dose of 60 milligrams was administered.

- (a) Circle the response below that best describes the type of relationship between M and t .

EXPONENTIAL GROWTH

EXPONENTIAL DECAY

- (b) Write down an equation for M in terms of t .

$$M = 60 e^{-0.12t} \quad //$$

- (c) Determine the mass of drug remaining in the bloodstream after one day.

$$\begin{aligned} m \Big|_{t=24} &= 60 e^{-0.12(24)} \\ &= \underline{3.37 \text{ mg}} \quad \checkmark \end{aligned}$$

- (d) Determine, to the nearest hour, the time taken for less than one percent of the initial dose to remain in the bloodstream of the patient.

$$\begin{aligned} 0.01 &= e^{-0.12t} \\ t &= 38.376 \text{ hrs.} \\ &\approx 38 \text{ hrs.} \quad // \end{aligned}$$

(e) At what rate is the mass of the drug in the bloodstream changing

(i) after 12 hours?


$$\begin{aligned}\frac{dm}{dt} &= -0.12 (60 e^{-0.12 \times 12}) \\ &= -1.706 \text{ mg/hr} \quad \checkmark\end{aligned}$$

(ii) when 25mg of the drug remains?

$$\begin{aligned}\frac{dm}{dt} &= -0.12 \times 25 \\ &= -3 \text{ mg/hr.} \quad \checkmark\end{aligned}$$

Q7 (3 marks)

A section of the graph of the function $y = 0.5 - \sin x$ is shown below. Calculate the **enclosed area** between the function stated and the x axis as shown in the diagram.



$$\begin{aligned}\int_{\frac{\pi}{6}}^{\frac{5\pi}{6}} |0.5 - \sin x| dx \\ = 8.338 \text{ units}^2 \quad \checkmark\end{aligned}$$

END OF SECTION 2