

**Mathematics Methods 3 & 4**  
**Test 1 2016**

Section 1 Calculator Free  
**Differentiation, Anti-differentiation and their applications.**

STUDENT'S NAME \_\_\_\_\_

MARKING KEY

DATE: Friday 4<sup>th</sup> March

TIME: 25 minutes

MARKS: 27

**INSTRUCTIONS:**

Standard Items: Pens, pencils, drawing templates, eraser, Formula sheet.

Questions or parts of questions worth more than 2 marks require working to be shown to receive full marks.

1. (6 marks)

Differentiate the following. Do not simplify your answer.

(a)  $y = (4x + 7)^3(9x - 5)$  let  $u = (4x + 7)^3$   $v = (9x - 5)$  [3]

$$u' = 3(4x + 7)^2(4) \quad v' = 9$$

$$= 12(4x + 7)^2$$

$$\frac{dy}{dx} = u'v + v'u$$

$$= 12(4x + 7)^2(9x - 5) + 9(4x + 7)^3 \quad \checkmark \checkmark \checkmark$$

(b)  $y = \frac{\sqrt{2x^5}}{\sqrt{x+7}}$  let  $u = (2x^5)^{\frac{1}{2}}$   $u' = 5x^4(2x^5)^{-\frac{1}{2}}$  [3]

$$v = (x+7)^{\frac{1}{2}} \quad v' = \frac{1}{2}(x+7)^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{u'v - v'u}{v^2}$$

$$= \frac{5x^4(2x^5)^{-\frac{1}{2}}(x+7)^{\frac{1}{2}} - \frac{1}{2}(x+7)^{-\frac{1}{2}}(2x^5)^{\frac{1}{2}}}{x+7} \quad \checkmark \checkmark$$

2. (3 marks)

Determine  $\int 2x(7-3x^2)^4 dx$

$$= \frac{2x(7-3x^2)^5}{5(6x)} + c$$

$$= \frac{(7-3x^2)^5}{-15} + c$$

3. (3 marks)

Given that  $\int_1^a (2x-3)dx = 6$ , determine  $a$ .

$$= [x^2 - 3x]_1^a$$

$$= (a^2 - 3a) - (1^2 - 3(1))$$

$$\therefore a^2 - 3a + 2 = 6$$

$$a^2 - 3a - 4 = 0$$

$$\therefore a = -1$$

$$a = 4$$

4. (6 marks)

The air in a hot air balloon is being inflated such that the rate of change of its volume at any time  $t$ , minutes, is given as:

$$\frac{dV}{dt} = 3t^2 - 2t \quad \text{for } t \geq 0$$

If initially the balloon has  $3 \text{ m}^3$  of air in it, determine:

(a) The rate of change in volume when  $t = 1$ . Explain the meaning of this. [2]

$$\begin{aligned} \frac{dV}{dt} &= 3(1) - 2(1) \\ &= 1 \text{ m}^3/\text{min} \quad \checkmark \end{aligned}$$

The balloon is instantaneously increasing in volume by  $1 \text{ m}^3/\text{min}$  at  $t = 1$ .  $\checkmark$

(b) For what values of  $t$  the volume is increasing. [2]

$$\frac{dV}{dt} > 0.$$

$$3t^2 - 2t > 0 \quad \checkmark \quad \therefore t > \frac{2}{3} \quad \checkmark$$

$$t(3t - 2) > 0$$

$$\begin{aligned} t &> 0, \quad t > \frac{2}{3} \\ t &< 0 \end{aligned}$$

(c) The volume of the balloon after five minutes. [2]


$$V(t) = t^3 - t^2 + 3 \quad \checkmark$$

$$V(5) = (5)^3 - (5)^2 + 3$$

$$= 103 \text{ m}^3 \quad \checkmark$$

5. (4 marks)

The graph of  $y = f(x)$  is shown below.



Given  $\int_{-3}^0 f(x) dx = 4$  and  $\int_0^2 f(x) dx = -1$ , determine the following:

(a)  $\int_{-3}^2 f(x) dx = 3$  ✓ [1]

(b)  $\int_0^2 5f(x) dx = 5 \int_0^2 f(x) dx$  [1]  
 $= 5(-1)$   
 $= -5$  ✓

(c)  $\int_{-3}^2 |f(x)| dx = 5$  ✓ [1]

(d) The area enclosed by  $f(x)$  and the  $x$  axis. [1]

$5 \text{ units}^2$  ✓

6. <sup>6</sup>/<sub>5</sub> marks)

Given the function  $y = (x+2)(x^2 - 4x + 4)$ .

(a) Determine the gradient of the tangent to the curve at  $x = 3$ . [3]

$$\begin{aligned}\frac{dy}{dx} &= 1(x^2 - 4x + 4) + (2x - 4)(x + 2) \checkmark \\ &= x^2 - 4x + 4 + (2x^2 - 8) \\ &= 3x^2 - 4x - 4 \checkmark \\ \frac{dy}{dx} \bigg|_{x=3} &= 3(9) - 4(3) - 4 \quad \text{gradient} = 11 \checkmark\end{aligned}$$

(b) Using calculus techniques, determine the nature of the stationary point at  $x = 2$ . [3]

$$\frac{d^2y}{dx^2} = 6x - 4 \quad \checkmark$$

$$\begin{aligned}\frac{d^2y}{dx^2} \bigg|_{x=2} &= 6(2) - 4 \\ &= 8 \quad \checkmark\end{aligned}$$

$\frac{d^2y}{dx^2}$  is positive  $\therefore$  minimum T.P.  $\checkmark$

**Mathematics Methods 3 & 4**  
**Test 1 2016**

Section 2 Calculator Assumed  
**Differentiation, Anti-differentiation and their applications.**

STUDENT'S NAME

MARKING KEY

DATE: Friday 4<sup>th</sup> March

TIME: 25 mins

MARKS: 23

**INSTRUCTIONS:**

Standard Items: Pens, pencils, drawing templates, eraser, Formula sheet.

Special Items: Three calculators, notes on one side of a single A4 page (these notes to be handed in with this assessment)

Questions or parts of questions worth more than 2 marks require working to be shown to receive full marks.

7. (6 marks)

The volume  $V \text{ cm}^3$  of water in a vessel is given by  $V = \frac{1}{6}\pi x^3$ , where  $x \text{ cm}$  is the depth of the water in the cylinder in cm.

- (a) Determine an approximation for the change in depth when the volume of water changes from 200 to 210  $\text{cm}^3$ . [3]

$$\begin{aligned}
 V &= 200 & \checkmark & \quad \frac{dV}{dx} = \frac{\pi x^2}{2} \\
 x &= \sqrt[3]{\frac{1200}{\pi}} \approx 7.26 & \checkmark & \quad \therefore \frac{dx}{dV} = \frac{2}{\pi x^2}
 \end{aligned}$$

$$\begin{aligned}
 \Delta x &= \frac{2}{\pi (7.26)^2} \times 10 & \checkmark & \quad \Delta x = 0.121 \text{ cm} \checkmark
 \end{aligned}$$

- (b) Determine the percentage change in the volume of the vessel if the depth has increased by 6%. [3]

$$\begin{aligned}
 \Delta V &= \frac{\pi x^2}{2} \times 0.06x & \quad \frac{\Delta V}{V} &= 0.18 \\
 &= 0.03\pi x^3 & \checkmark & \quad = 18\% \checkmark
 \end{aligned}$$

$$\frac{\Delta V}{V} = \frac{0.03\pi x^3}{\frac{1}{6}\pi x^3} \checkmark$$

8. (4 marks)

A company manufacturing a new bike determines that the marginal cost (in dollars) for the production of the  $n^{\text{th}}$  unit is given by the expression:

$$\frac{dC}{dn} = \frac{200000}{(n+20)^2}$$

- (a) The initial set up cost is \$ 10 000 (i.e. the cost of producing no bikes is \$ 10 000). Show that the expression for the total cost of producing  $n$  bikes is:

$$C = \frac{-200000}{n+20} + 20000$$

$$C(0) = 10000 \quad [2]$$

$$C = \int \frac{dC}{dn}$$

$$10000 = \frac{-200000}{20} + C$$

$$= \frac{-200000}{n+20} + C \quad \checkmark$$

$$C = 20000 \quad \checkmark$$

$$\therefore C = \frac{-200000}{n+20} + 20000$$

- (b) If the company sells each bike for \$200, how many bikes must be sold before it first makes a profit? [2]


$$\text{Solve } 200n = \frac{-200000}{n+20} + 20000 \quad \checkmark$$

$$n = 90.99$$

The company must sell 91 bikes before making a profit.  $\checkmark$

9. (7 marks)

A man launches his boat from point A on a bank of a straight river, 3 km wide, and wants to reach point B, 8 km downstream on the opposite bank, as quickly as possible.



He could proceed in any of three ways:

1. Row his boat directly across the river to point C and then run to B
2. Row directly to B
3. Row to some point D between C and B and then run to B

(a) Given that  $time = \frac{distance}{speed}$  and  $x$  is the distance from C to D, show that the time ( $t$ ) taken for the man to travel from A to B can be represented by the equation. [2]

$$t = \frac{\sqrt{x^2 + 9}}{6} + \frac{8 - x}{8}$$

$$t = \frac{\overline{AD}}{6} + \frac{\overline{DB}}{8}$$

$$\therefore t = \frac{\sqrt{x^2 + 9}}{6} + \frac{8 - x}{8}$$

Given  $x = \overline{CD}$

$$\overline{AD} = \sqrt{3^2 + x^2}$$

$$\overline{AD} = \sqrt{x^2 + 9}$$

$$\overline{DB} = 8 - \overline{CD}$$

$$\overline{DB} = 8 - x$$





- (b) Using calculus techniques, determine the minimum time taken by the man to reach point B and the distance he would travel by foot to achieve this minimum time. [5]

$$t(x) = \frac{1}{6}(x^2+9)^{\frac{1}{2}} + \frac{1}{8}(8-x)$$

$$t'(x) = \frac{1}{12}(x^2+9)^{-\frac{1}{2}} \cdot (2x) - \frac{1}{8}$$

$$t'(x) = \frac{x}{6(x^2+9)^{\frac{1}{2}}} - \frac{1}{8} \quad \checkmark \quad 0 = \frac{4x - 3\sqrt{x^2+9}}{24\sqrt{x^2+9}}$$

Solve  $t'(x) = 0$

$$x = \frac{9\sqrt{7}}{7}$$

$$x = \frac{9}{\sqrt{7}} \text{ via CP. } \checkmark \quad (3.40)$$

Verify  $\frac{9}{\sqrt{7}}$  is a min.

$$t''\left(\frac{9}{\sqrt{7}}\right) = \frac{7\sqrt{7}}{1152} > 0 \quad \therefore \frac{9}{\sqrt{7}} \text{ is a min. } \checkmark$$

(0.0161)

$$\begin{aligned} \text{time taken} &= \frac{1}{6}\left[\left(\frac{9}{\sqrt{7}}\right)^2 + 9\right]^{\frac{1}{2}} + \frac{1}{8}\left(8 - \frac{9}{\sqrt{7}}\right) \\ &= \frac{2}{\sqrt{7}} + 0.5748 \\ &= 1.3307 \end{aligned}$$

$$\begin{aligned} \text{Running dist} &= 8 - \frac{9}{\sqrt{7}} \\ &= 4.598 \text{ km} \end{aligned}$$

It would take the man 1.3307 hours to reach B. He would run 4.598 km.  $\checkmark$

10. (6 marks)

A particle moves in rectilinear motion with a velocity of 7 m/s as it passes through a fixed point O.

$t$  is the number of seconds since passing through O. Acceleration  $a$  is defined as  $a = mt - n$ , where  $m$  and  $n$  are constants.

When  $t = 1$ , the velocity is 12 m/s, and when  $t = 7$  the particle is instantaneously at rest.

(a) Calculate the values of  $m$  and  $n$ .

[3]

$$V = \frac{mt^2}{2} - tn + c$$

$$V(0) = 7 \quad \therefore c = 7$$

$$\therefore V(t) = \frac{mt^2}{2} - tn + 7 \quad \checkmark$$

$$\underline{t=1 \quad V=12}$$

$$12 = \frac{m}{2} - n + 7 \quad \checkmark$$

$$m = -2 \quad \checkmark$$

$$n = -6 \quad \checkmark$$

$$\underline{t=7 \quad V=0}$$

$$0 = \frac{49m}{2} - 7n + 7$$

(b) Hence, determine the expression for the velocity as a function of time.

[1]

$$V(t) = -t^2 + 6t + 7 \quad \checkmark$$

(c) Determine when and where the maximum velocity is attained.

[2]

max  $V$  occurs when  $\frac{dv}{dt} = 0$

$$\frac{dv}{dt} = -2t + 6$$

$$0 = -2t + 6$$

$$t = 3 \quad \checkmark$$

$$x(t) = -\frac{1}{3}t^3 + 3t^2 + 7t$$

$$x(3) = 39 \quad \checkmark$$