

**Mathematics Methods Units 3/4
Test 2 2017**

Section 1 Calculator Free
Applications of Calculus

STUDENT'S NAME _____

DATE: Tuesday 28 March

TIME: 25 minutes

MARKS: 27

INSTRUCTIONS:

Standard Items: Pens, pencils, drawing templates, eraser

Questions or parts of questions worth more than 2 marks require working to be shown to receive full marks.

1. (9 marks)

Differentiate each of the following with respect to x . (Do not simplify your answers):

(a) $y = x^5 e^{-3x}$ [2]

(b) $y = \cos\left(\sqrt{7 + e^x}\right)$ [3]

(c) $y = f(5 - 3x)$ where f is a function [2]

(d) $y = \int_x^1 (1 + 2t)^2 dt$ [2]

2. (9 marks)

(a) Determine:


(i) $\int 2x + e^{-2x} + e \, dx$ [3]

(ii) $\int \frac{xe^{1-2x^2}}{2} \, dx$ [3]

(b) Evaluate $\int_1^{\pi} \frac{d}{dx} \left(\frac{\sin x}{x^2 + 1} \right) dx$ [3]

3. (5 marks)

Calculate the area enclosed between the functions $e^x - 1$, $x(x-2)$ and the line $x = 2$ as indicated on the graph below:



4. (4 marks)

A continuous function $f(x)$ is increasing on the interval $0 < x < 2$ and decreasing on the interval $2 < x < 5$. Some of its values are given in the table below:

x	0	1	2	3	4	5
$f(x)$	5	17	24	13	0	-29

The function $F(x)$ is defined, for $0 \leq x \leq 5$, by $F(x) = \int_0^x f(t) dt$.

(a) At which value of x in the interval $0 \leq x \leq 5$ is $F(x)$ greatest? Justify your answer.

[2]

(b) At which value of x in the interval $0 \leq x \leq 5$ is $F'(x)$ greatest? Justify your answer.

[2]

**Mathematics Methods Units 3/4
Test 2 2017**

Section 2 Calculator Assumed
Applications of Calculus

STUDENT'S NAME _____

DATE: Tuesday 28 March

TIME: 25 minutes

MARKS: 25

INSTRUCTIONS:

Standard Items: Pens, pencils, drawing templates, eraser

Special Items: Three calculators, notes on one side of a single A4 page (these notes to be handed in with this assessment)

Questions or parts of questions worth more than 2 marks require working to be shown to receive full marks.

5. (5 marks)

During a volcanic eruption a rock is ejected from the top of the volcano. The rock rises upward and then falls onto a flat plain 1500 metres below the top of the volcano. During its flight, the vertical velocity of the rock, v m/s, is given by

$$v = 160 - 9.8t$$

Where t seconds is the time after the ejection of the rock

(a) How high does the rock rise above the top of the volcano? [3]

(b) How long does it take for the rock to reach the plain below? [2]

6. (6 marks)

A radioactive substance is decaying exponentially, according to the formula

$$A(t) = A_0 e^{-kt}, \text{ where } A(t) \text{ kg is the amount at time } t \text{ years.}$$

- (a) Determine k , correct to 4 decimal places, given that the half-life of the substance is 12 years. [2]

A second radioactive substance is also decaying exponentially, according to the formula

$$B(t) = B_0 e^{-0.04t}, \text{ where } B(t) \text{ kg is the amount at time } t \text{ years.}$$

- (b) Which of these substances is decaying faster? Justify your answer briefly. [1]

At a certain location there was exactly the same amount of these two substances at the beginning of the year 2017.

- (c) In what year will the ratio of the amount of one of these substances to the other be 2:1? [3]

7. (7 marks)

The rate of population change of a bacteria culture is modelled by $\frac{dP}{dt} = 100e^{-0.01t}$ where t is in hours.

(a) Determine the initial instantaneous rate of change of P with respect to t . [1]

(b) Describe the rate of change for large values of t . [1]

(b) Determine the net change in population during the first 10 hours. [2]

(c) Determine the average change in population during the first 10 hours. [1]

(d) Given that the initial population was 100, determine the maximum population size. Show clearly how you obtained your answer. [2]

8. (7 marks)

The acceleration, $a(t) \text{ m s}^{-2}$, of an object moving in a straight line is given by:

$$a(t) = At + B, \text{ where } A \text{ and } B \text{ are non-zero constants.}$$

The object is at rest initially and again after 10 seconds, and the object returns to its initial position after T seconds.

(a) Evaluate T [4]

(b) Evaluate A and B given that the acceleration is positive initially and that the object travels a distance of 1 kilometre in the first T seconds. [3]