



Mathematics Methods U3/4
Test 3 2022

Section 1 Calculator Free
Discrete Random Variables

STUDENT'S NAME _____

DATE: Monday 9th May

TIME: 15 minutes

MARKS: 14

INSTRUCTIONS:


Standard Items: Pens, pencils, drawing templates, eraser

Questions or parts of questions worth more than 2 marks require working to be shown to receive full marks.

This page has been left intentionally blank

1. (7 marks)

The discrete random variable X can take the values 1, 2, 3, 4, 5. The probability distribution for X is shown graphically below.



Determine:

(a) the value of k . [2]

(b) $P(X \geq 3)$ [1]

(c) $P(X = 3 | X \geq 3)$ [2]

(d) the expected value. [2]

2. (7 marks)

(a) A discrete random variable X , where $X = 0, 1, 2, 3, 4, 5, 6, 7$ has a uniform distribution.

(i) Determine the expected value and variance of X . [3]

(ii) A discrete random variable is defined by $Y = 5 - 2X$. Calculate the expected value and variance of Y . [2]

(b) A discrete uniform distribution Z , has outcomes 0 to n and have an expected value, $E(X) = 5.5$. Calculate the value of n . [2]



**Mathematics Methods U3/4
Test 3 2022**

**Section 2 Calculator Assumed
Discrete Random Variables**

STUDENT'S NAME _____

DATE: Monday 9th May

TIME: 35 minutes

MARKS: 34

INSTRUCTIONS:

Standard Items: Pens, pencils, drawing templates, eraser

Special Items: Three calculators, notes on one side of a single A4 page (these notes to be handed in with this assessment)

Questions or parts of questions worth more than 2 marks require working to be shown to receive full marks.

This page has been left intentionally blank

3. (6 marks)

Tokens numbered 1 to 20 are placed in a bag, and one is selected at random:

- Let $X = 1$ if a prime number is selected, and $X = 0$ otherwise.
- Let $Y = 1$ if a number greater than 8 is selected, and $Y = 0$ otherwise.

(a) Determine the probability of success of X and Y respectively. [2]

(b) Calculate the mean and variance of X . [1]

(c) Calculate the mean and variance of Y . [1]

(d) Compare the standard deviation of X and Y and justify why this occurred? [2]

4. (9 marks)

It is known the probability of a bread roll pen being below the satisfactory weight for sale in a large batch is 0.12. At the bakery, bread rolls are sold in packets of 6.

(a) Describe the probability distribution function. [2]

(b) Determine $E(X)$ and $\text{st.dev}(X)$ [2]

(c) Determine the probability that

(i) A randomly selected packet has greater than 2 bread rolls that are below the satisfactory weight for sale. [1]

(ii) that there were no more than 1 bread roll that is below the satisfactory weight for sale in each packet if 6 packets were randomly selected. [2]

(iii) in a large order of 20 packets of bread rolls that no more than 3 of these have greater than 2 bread rolls that are below the satisfactory weight for sale. [2]

5. (10 marks)

A slot machine is programmed to operate at random, making various payouts after patrons pay \$2 and press a start button. The random variable X is the amount of the payout, in dollars, in one play of the machine. Each payout can be assumed to be independent of other payouts.

The probability, P , that the machine makes a certain payout, x , is shown in the table below.

Payout (\$) x	0	1	2	5	10	20	50	100
Probability $P(X = x)$	0.25	0.45	0.2125	0.0625	0.0125	0.005	0.005	0.0025

(a) Determine the probability that

(i) in one play of the machine, a payout of more than \$1 is made. [1]

(ii) in ten plays of the machine, it makes a payout of \$5 no more than once. [2]

(iii) in five plays of the machine, the second payout of \$1 occurs on the fifth play. [3]

(b) Calculate the mean and standard deviation of X . [2]

(c) In the long run, what percentage of the patron's money is returned to them? [2]

6. (9 marks)

In the mailroom of a large company, it is known that 20% of incoming letters contain an invoice. Let X be the number of randomly chosen letters that are opened until an invoice is discovered.

(a) Complete the table below for the values of $x = 1, 2, 3,$ and 4 . [2]

x	1	2	3	4
$P(X = x)$				

(b) Determine the rule for $P(X = x)$ for any integer value greater than 0. [2]

(c) Calculate

(i) $P(X = 10)$ [1]

(ii) $P(3 \leq X \leq 6)$ [2]

(iii) the smallest value of k , so that $P(X = k) < 0.001$. [2]