

# Physics ATAR - Year 11

## Thermal Physics Validation Test 2018

Name:
-------

Mark:	/ 47
=	%

Time Allowed: 50 Minutes

Notes to Students:

1. You must include **all** working to be awarded full marks for a question.
2. Marks will be deducted for incorrect or absent units and answers stated to an incorrect number of significant figures.
3. **No** graphics calculators are permitted – scientific calculators only.

### ADDITIONAL FORMULAE AND DATA

- Triple Point of Water = 0.01°C = 273.16 K
- $\frac{T_{\theta}}{100} = \frac{X_{\theta} - X_0}{X_{100} - X_0} = \frac{R_{\theta} - R_0}{R_{100} - R_0}$
- $K = C + 273.15$

Question 1

(2 marks)

The Celsius scale was originally calibrated as a centigrade scale with 0 °C for the freezing point of water and 100 °C for the boiling point of water at 1 atm pressure as the fixed points. These were later deemed unreliable and changed to two other fixed points. State these fixed points and explain why they are considered more reliable.

---

---

---

---

**Question 2****(3 marks)**

A pan of water is heated from 25.0 °C to 355 K. Calculate the change in temperature in the Kelvin scale.

**Question 3****(3 marks)**

The electrical resistance an uncalibrated thermostat thermometer is 0.250  $\Omega$  in ice water and 3.56  $\Omega$  in boiling water. When placed in a liquid with an unknown temperature the resistance is recorded as 3.99  $\Omega$ . Calculate the value of the unknown temperature.

**Question 4****(3 marks)**

Calculate the amount of energy the needs to be removed from a 1.55 kg piece of metal to reduce

**Question 5**

**(4 marks)**

Metal A has a specific heat capacity of 2.00x. Metal B has a specific heat of 4.50x. Equal masses of each metal are provided with equal amounts of heat and Metal A records a  $\Delta T$  of 5.55 K. Calculate, through an adequate mathematical relationship, the  $\Delta T$  of Metal B.

**Question 6**

**(8 marks)**

A large bucket of water at 30.0 °C sits next to a small cup of water at 30.0 °C.



- (a) Choose which volume of water has more internal energy (Circle your chosen answer) (1 mark)
- (i) The bucket
  - (ii) The cup
  - (iii) They both have the same amount of internal energy

(b) Explain your answer to (a). (3 marks)

---

---

---

---

---

- (c) Choose which has the fastest moving molecules (Circle your chosen answer) (1 mark)
- (i) The bucket
  - (ii) The cup
  - (iii) They both have the same mean speed of particles

(d) Explain your answer to (c). (3 marks)

---

---

---

---

---

**Question 7** (4 marks)

Calculate the total energy required to vaporise  $5.00 \times 10^2$  grams of water initially at  $25.0 \text{ }^\circ\text{C}$ .

**Question 8****(7 marks)**

A  $1.50 \times 10^3 \text{ W}$  copper kettle of mass  $0.350 \text{ kg}$  contains  $1.80 \text{ kg}$  of water at  $25.0 \text{ }^\circ\text{C}$ .

( $c_{\text{Cu}} = 390 \text{ Jkg}^{-1}\text{K}^{-1}$ )

( $c_{\text{w}} = 4180 \text{ Jkg}^{-1}\text{K}^{-1}$ )

- (a) Calculate the amount of energy required to bring the water (and the copper) to its boiling point.

**(3 marks)**

- (b) Calculate the time, in minutes, required to bring the water to boil

**(4 marks)****Question 9****(8 marks)**

A mass of aluminium at  $90.0 \text{ }^\circ\text{C}$  is immersed in an insulated  $0.255 \text{ kg}$  volume of water initially at  $20.0 \text{ }^\circ\text{C}$ . The final temperature of the mixture is measured to be  $24.4 \text{ }^\circ\text{C}$ . ( $c_{\text{Al}} = 900.0 \text{ Jkg}^{-1}\text{K}^{-1}$ )

- (a) Calculate the mass of aluminium required to produce the final temperature.

**(4 marks)**

(b) State one assumption made in the above calculation.

(1 mark)

---

---

(c) In reality, state and explain whether the required mass of aluminium would be greater, less, or no difference to raise the temperature of the mixture to 24.4 °C.

(3 marks)

---

---

---

---

**Question 10**

**(5 marks)**

1.40 kg of water is placed in a calorimeter of mass 1.00 kg and specific heat capacity of  $655 \text{ Jkg}^{-1}\text{K}^{-1}$ . Both are cooled to  $4.00^\circ\text{C}$ . 0.250 kg of copper at  $90.0^\circ\text{C}$  is then placed into the calorimeter. Calculate the resulting temperature of the mixture.

( $c_{\text{Cu}} = 390 \text{ Jkg}^{-1}\text{K}^{-1}$ )

( $c_{\text{w}} = 4180 \text{ Jkg}^{-1}\text{K}^{-1}$ )

**END OF TEST**