



MATHEMATICS: SPECIALIST

UNITS 3C AND 3D

FORMULA SHEET 2015

Copyright

© School Curriculum and Standards Authority, 2015

This document—apart from any third party copyright material contained in it—may be freely copied, or communicated on an intranet, for non-commercial purposes by educational institutions, provided that it is not changed in any way and that the School Curriculum and Standards Authority is acknowledged as the copyright owner.

Copying or communication for any other purpose can be done only within the terms of the Copyright Act or by permission of the Authority.

Copying or communication of any third party copyright material contained in this document can be done only within the terms of the Copyright Act or by permission of the copyright owners.

This document is valid for teaching and examining until 31 December 2015.

Index

Vectors **3**

Trigonometry **3**

Functions **4**

Matrices **5**

Complex numbers **6**

Exponentials and logarithms **7**

Mathematical reasoning **7**

Measurement **8**

Vectors

Magnitude:	$ (a_1, a_2, a_3) = \sqrt{a_1^2 + a_2^2 + a_3^2}$	
Dot product:	$\mathbf{a} \cdot \mathbf{b} = \mathbf{a} \mathbf{b} \cos \theta = a_1b_1 + a_2b_2 + a_3b_3$	
Triangle inequality:	$ \mathbf{a} + \mathbf{b} \leq \mathbf{a} + \mathbf{b} $	
Vector equation of a line in space:	one point and the slope:	$\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$
	two points A and B:	$\mathbf{r} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$
Cartesian equations of a line in space:	$\frac{x - a_1}{b_1} = \frac{y - a_2}{b_2} = \frac{z - a_3}{b_3}$	

Parametric form of vector equation of a line in space:

$$\begin{aligned} x &= a_1 + \lambda b_1, \dots (1) \\ y &= a_2 + \lambda b_2, \dots (2) \\ z &= a_3 + \lambda b_3, \dots (3) \end{aligned}$$

Vector equation of a plane in space:	$\mathbf{r} \cdot \mathbf{n} = c$	or	$\mathbf{r} = \mathbf{a} + \lambda \mathbf{b} + \mu \mathbf{c}$
--------------------------------------	-----------------------------------	----	---

Trigonometry

In any triangle ABC :	$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
	$a^2 = b^2 + c^2 - 2bc \cos A$
	$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$
	$\text{Area} = \frac{1}{2} ab \sin C$

In a circle of radius r , for an arc subtending angle θ (radians) at the centre:

Length of arc	$= r\theta$	
Area of segment	$= \frac{1}{2} r^2 (\theta - \sin \theta)$	Area of sector
		$= \frac{1}{2} r^2 \theta$

Identities:	$\cos^2 \theta + \sin^2 \theta = 1$	$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$
	$\cos (\theta \pm \varphi) = \cos \theta \cos \varphi \mp \sin \theta \sin \varphi$	$= 2\cos^2 \theta - 1$
		$= 1 - 2\sin^2 \theta$
	$\sin (\theta \pm \varphi) = \sin \theta \cos \varphi \pm \cos \theta \sin \varphi$	$\sin 2\theta = 2 \sin \theta \cos \theta$
	$\tan (\theta \pm \varphi) = \frac{\tan \theta \pm \tan \varphi}{1 \mp \tan \theta \tan \varphi}$	$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$
	$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$	$\lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0$

Simple Harmonic Motion: If $\frac{d^2x}{dt^2} = -k^2x$ then $x = A \sin(kt + \alpha)$ or $x = A \cos(kt + \beta)$ and

$v^2 = k^2 (A^2 - x^2)$, where A is the amplitude of the motion, α and β are phase angles, v is the velocity and x is the displacement.

Functions

Differentiation: If $f(x) = y$ then $f'(x) = \frac{dy}{dx}$ If $f(x) = x^n$ then $f'(x) = nx^{n-1}$

 If $f(x) = e^x$ then $f'(x) = e^x$ If $f(x) = \ln x$ then $f'(x) = \frac{1}{x}$

 If $f(x) = \sin x$ then $f'(x) = \cos x$ If $f(x) = \cos x$ then $f'(x) = -\sin x$

 If $f(x) = \tan x$ then $f'(x) = \sec^2 x = \frac{1}{\cos^2 x}$

Product rule: If $y = f(x) g(x)$ or If $y = uv$

 then $y' = f'(x) g(x) + f(x) g'(x)$ then $\frac{dy}{dx} = \frac{du}{dx} v + u \frac{dv}{dx}$

Quotient rule: If $y = \frac{f(x)}{g(x)}$ or If $y = \frac{u}{v}$

 then $y' = \frac{f'(x) g(x) - f(x) g'(x)}{(g(x))^2}$ then $\frac{dy}{dx} = \frac{\frac{du}{dx} v - u \frac{dv}{dx}}{v^2}$

Incremental formula: $\delta y \approx \frac{dy}{dx} \delta x$ or $f(x+h) - f(x) \approx f'(x)h$

Chain rule: If $y = f(g(x))$

 then $y' = f'(g(x)) g'(x)$ or If $y = f(u)$ and $u = g(x)$

 then $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$

Integration:

Powers: $\int x^n dx = \frac{x^{n+1}}{n+1} + c, n \neq -1$

Exponentials: $\int e^x dx = e^x + c$ Logarithms: $\int \frac{1}{x} dx = \ln|x| + c$

Trigonometric: $\int \sin x dx = -\cos x + c$
 $\int \cos x dx = \sin x + c$
 $\int \frac{1}{\cos^2 x} dx = \tan x + c$

Fundamental Theorem of Calculus:

$$\frac{d}{dx} \int_a^x f(t) dt = f(x) \quad \text{and} \quad \int_a^b f'(x) dx = f(b) - f(a)$$

See next page

Functions

Quadratic function:

$$\text{If } y = ax^2 + bx + c \text{ and } y = 0, \text{ then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \text{ for } x \in \mathbb{C}$$

Piecewise-defined functions:

$$\text{Absolute value function: } |x| = \begin{cases} x, & \text{for } x \geq 0 \\ -x, & \text{for } x < 0 \end{cases}$$

$$\text{Sign function: } \operatorname{sgn}(x) = \begin{cases} 1, & \text{for } x > 0 \\ 0, & \text{for } x = 0 \\ -1, & \text{for } x < 0 \end{cases}$$

Greatest integer function: $\operatorname{int}(x) = \text{greatest integer } \leq x \text{ for all } x$ **Matrices**

$$\text{If } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \text{ then } |A| = \det A = ad - bc$$

$$A^{-1} = \frac{1}{(ad - bc)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\text{Dilation} = \begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix}$$

$$\text{Shear} = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \text{ or } \begin{bmatrix} 1 & 0 \\ a & 1 \end{bmatrix}$$

$$\text{Rotation} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$\text{Reflection} = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix}$$

Complex numbers

For $z = a + ib$, where $i^2 = -1$

Argument: $\arg z = \theta$, where $\tan \theta = \frac{b}{a}$ and $-\pi < \theta \leq \pi$

Modulus: $\text{mod } z = |z| = |a + ib| = \sqrt{a^2 + b^2} = r$

Product: $|z_1 z_2| = |z_1| |z_2|$ $\arg(z_1 z_2) = \arg z_1 + \arg z_2$

Quotient: $\frac{|z_1|}{|z_2|} = \frac{|z_1|}{|z_2|}$ $\arg \frac{z_1}{z_2} = \arg z_1 - \arg z_2$

Polar form:

For $z = r \text{ cis } \theta$, where $r = |z|$ and $\theta = \arg z$:

$$\text{cis}(\theta + \varphi) = \text{cis } \theta \text{ cis } \varphi$$

$$\text{cis}(-\theta) = \frac{1}{\text{cis } \theta}$$

$$z_1 z_2 = r_1 r_2 \text{ cis } (\theta + \varphi)$$

$$\text{cis } \theta = \cos \theta + i \sin \theta$$

$$\text{cis}(0) = 1$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} \text{ cis } (\theta - \varphi)$$

Exponential form:

$$z = r e^{i\theta}, \text{ where } r = |z| \text{ and } \theta = \arg z$$

For complex conjugates:

$$z = a + bi$$

$$z = r \text{ cis } \theta$$

$$z = r e^{i\theta}$$

$$z \bar{z} = |z|^2$$

$$\overline{z_1 + z_2} = \bar{z}_1 + \bar{z}_2$$

$$\bar{z} = a - bi$$

$$\bar{z} = r \text{ cis } (-\theta)$$

$$\bar{z} = r e^{-i\theta}$$

$$z^{-1} = \frac{1}{z} = \frac{\bar{z}}{|z|^2}$$

$$\overline{z_1 z_2} = \bar{z}_1 \bar{z}_2$$

Exponentials and logarithms

For $a, b > 0$ and m, n real:

$$a^m a^n = a^{m+n}$$

$$a^0 = 1$$

$$(a^m)^n = a^{mn}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

$$a^{-n} = \frac{1}{a^n}$$

$$(ab)^m = a^m b^m$$

For m an integer and n a positive integer:

$$a^{\frac{1}{n}} = \sqrt[n]{a}$$

$$a^{\frac{m}{n}} = \sqrt[n]{a^m} = (\sqrt[n]{a})^m$$

For a, b, y, m and n positive real and k real:

$$1 = a^0 \Leftrightarrow \log_a 1 = 0$$

$$\log_a mn = \log_a m + \log_a n$$

$$\log_a m = \frac{\log_b m}{\log_b a} \quad (\text{change of base})$$

$$y = a^x \Leftrightarrow \log_a y = x$$

$$a = a^1 \Leftrightarrow \log_a a = 1$$

$$\log_a (m^k) = k \log_a m$$

If $\frac{dP}{dt} = kP$, then $P = P_0 e^{kt}$ **Mathematical reasoning**

De Moivre's theorem:

$$(\text{cis } \theta)^n = (\cos \theta + i \sin \theta)^n$$

$$(\text{cis } \theta)^n = \cos n\theta + i \sin n\theta$$

$$z^n = |z|^n \text{cis } (n\theta)$$

$$z^{\frac{1}{q}} = |z|^{\frac{1}{q}} \left[\cos \left[\frac{\theta + 2\pi k}{q} \right] + i \sin \left[\frac{\theta + 2\pi k}{q} \right] \right] \text{ for } k \text{ an integer.}$$

Measurement

Circle: $C = 2\pi r = \pi D$, where C is the circumference,
 r is the radius and D is the diameter
 $A = \pi r^2$, where A is the area

Triangle: $A = \frac{1}{2}bh$, where b is the base and h is the perpendicular height

Parallelogram: $A = bh$

Trapezium: $A = \frac{1}{2}(a + b)h$, where a and b are the lengths of the parallel sides

Prism: $V = Ah$, where V is the volume and A is the area of the base

Pyramid: $V = \frac{1}{3} Ah$

Cylinder: $S = 2\pi rh + 2\pi r^2$, where S is the total surface area
 $V = \pi r^2 h$

Cone: $S = \pi rs + \pi r^2$, where s is the slant height
 $V = \frac{1}{3} \pi r^2 h$

Sphere: $S = 4\pi r^2$
 $V = \frac{4}{3} \pi r^3$

Note: Any additional formulas identified by the examination panel as necessary will be included in the body of the particular question.