



# PERTH MODERN SCHOOL

Exceptional schooling. Exceptional students.

INDEPENDENT PUBLIC SCHOOL

WAEP Semester One Examination, 2019

Question/Answer booklet

## MATHEMATICS SPECIALIST UNIT 1

Section Two:

Calculator-assumed

# SOLUTIONS

Student number: In figures

--	--	--	--	--	--	--	--

In words

---

Your name

---

### Time allowed for this section

Reading time before commencing work: ten minutes

Working time: one hundred minutes

### Materials required/recommended for this section

#### *To be provided by the supervisor*

This Question/Answer booklet

Formula sheet (retained from Section One)

#### *To be provided by the candidate*

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: drawing instruments, templates, notes on two unfolded sheets of A4 paper, and up to three calculators approved for use in this examination

### Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

## Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
<b>Total</b>					100

## Instructions to candidates

1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet preferably using a blue/black pen. Do not use erasable or gel pens.
3. You must be careful to confine your answer to the specific question asked and to follow any instructions that are specified to a particular question.
4. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
5. It is recommended that you do not use pencil, except in diagrams.
6. Supplementary pages for planning/continuing your answers to questions are provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
7. The Formula sheet is not to be handed in with your Question/Answer booklet.

Section Two: Calculator-assumed

65% (98 Marks)


This section has **thirteen (13)** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time: 100 minutes.

**Question 9**

**(5 marks)**

In the diagram below,  $M$  is the midpoint of  $QR$ .



If  $\vec{OP} = \mathbf{p}$ ,  $\vec{OQ} = \mathbf{q}$  and  $\vec{OR} = \mathbf{r}$ , express the following in terms of  $\mathbf{p}$ ,  $\mathbf{q}$  and  $\mathbf{r}$ .

(a)  $\vec{PR}$ .

Solution
$\vec{PR} = \mathbf{r} - \mathbf{p}$
Specific behaviours
✓ correct expression

(1 mark)

(b)  $\vec{OM}$ .

Solution
$\vec{OM} = \vec{OQ} + \frac{1}{2}\vec{QR}$ $= \mathbf{q} + \frac{1}{2}(\mathbf{r} - \mathbf{q})$ $= \frac{1}{2}\mathbf{q} + \frac{1}{2}\mathbf{r}$
Specific behaviours
✓ indicates correct method ✓ correct expression

(2 marks)

(c)  $6\vec{MP}$ .

Solution
$\vec{MP} = \vec{MO} + \vec{OP} = \mathbf{p} - \frac{1}{2}\mathbf{q} - \frac{1}{2}\mathbf{r}$ $\vec{MP} = 6\mathbf{p} - 3\mathbf{q} - 3\mathbf{r}$
Specific behaviours
✓ indicates $\vec{MP}$ ✓ correct expression

(2 marks)

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

## Question 10

(8 marks)

Points  $P$ ,  $Q$  and  $R$  have coordinates  $(-2, 11)$ ,  $(8, 15)$  and  $(17, 3)$  respectively. Determine

(a)  $\overrightarrow{PQ}$ .

(1 mark)

Solution
$\overrightarrow{PQ} = (8, 15) - (-2, 11)$ $= (10, 4)$
Specific behaviours
✓ correct vector

(b)  $|\overrightarrow{QR}|$ .

(2 marks)

Solution
$\overrightarrow{QR} = (17, 3) - (8, 15)$ $= (9, -12)$
$ \overrightarrow{QR}  = 15$
Specific behaviours
✓ correct vector
✓ magnitude

(c)  $2\overrightarrow{PQ} - 60\mathbf{u}$ , where  $\mathbf{u}$  is a unit vector in the direction  $\overrightarrow{QR}$ .

(3 marks)

Solution
$\mathbf{u} = \frac{1}{15}(9, -12)$
$2\overrightarrow{PQ} - 60\mathbf{u} = 2(10, 4) - \frac{60}{15}(9, -12)$ $= (-16, 56)$
Specific behaviours
✓ indicates unit vector
✓ expression for result
✓ correct vector

(d) The coordinates of point  $S$ , given that  $\overrightarrow{RS} = \overrightarrow{QP}$ .


(2 marks)

Solution
$\overrightarrow{OS} = \overrightarrow{OR} + \overrightarrow{RS}$ $= \overrightarrow{OR} - \overrightarrow{PQ}$ $= (21, 3) - (10, 4)$ $= (7, -1)$
Specific behaviours
✓ expression for result
✓ correct coordinates

Question 11

(7 marks)

- (a) In the diagram below (not drawn to scale)  $A, B, C$  and  $D$  lie on a circle and  $EB$  and  $ED$  are tangents to the circle. If  $\angle BED = 54^\circ$  and  $\angle CDB = 20^\circ$ , determine the size of  $\angle CBD$ . (3 marks)



Solution
$\angle BDE = (180 - 54) \div 2 = 63$
$\angle CDE = 63 - 20 = 43$
$\angle CBD = \angle CDE = 43^\circ$ (AltSegment)
Specific behaviours
<ul style="list-style-type: none"> <li>✓ <math>\angle BDE</math></li> <li>✓ <math>\angle CDE</math></li> <li>✓ <math>\angle CBD</math></li> </ul>

- (b) Quadrilateral  $ABCD$  is such that  $CB = CD$ ,  $\angle BAD = 96^\circ$  and  $\angle BDC = 48^\circ$ .

- (i) Sketch a diagram to show this information. (1 mark)

Solution
Specific behaviours
✓ correct diagram

- (ii) Show that  $ABCD$  is cyclic and hence determine the size of  $\angle CAD$ . (3 marks)

Solution
$\angle CBD = \angle CDB = 48$ $\angle BCD = 180 - 2 \times 48 = 84$  $\angle BAD + \angle BCD = 96 + 84 = 180$ Hence cyclic as opposite angles supplementary.  $\angle CAD = \angle CBD = 48^\circ$ (Same arc)
Specific behaviours
<ul style="list-style-type: none"> <li>✓ use isosceles triangle for <math>\angle BCD</math></li> <li>✓ uses supplementary angles for cyclic</li> <li>✓ correct size of <math>\angle CAD</math></li> </ul>

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

## Question 12

(8 marks)

- (a) Show that the vectors
- $(8, -5)$
- and
- $(2.5, 4)$
- are perpendicular.

(2 marks)

Solution
$\begin{pmatrix} 8 \\ -5 \end{pmatrix} \cdot \begin{pmatrix} 2.5 \\ 4 \end{pmatrix} = 20 - 20 = 0$
Hence perpendicular as scalar (dot) product is 0.
Specific behaviours
<ul style="list-style-type: none"> <li>✓ uses dot product</li> <li>✓ explains result</li> </ul>

- (b) Determine, to the nearest degree, the angle between the vectors
- $(3, -2)$
- and
- $(-2, -4)$
- .

(2 marks)

Solution
Using CAS: $\theta = 82.87 \approx 83^\circ$
Or: $\theta = \cos^{-1}\left(\frac{2}{\sqrt{13} \times 2\sqrt{5}}\right)$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ indicates method</li> <li>✓ correct angle</li> </ul>

- (c) The vectors
- $(a, 2a + 3)$
- and
- $(a + 3, -2)$
- are perpendicular, where
- $a$
- is a constant. Determine the value(s) of
- $a$
- and the corresponding pair(s) of vectors.


(4 marks)

Solution
$\begin{pmatrix} a \\ 2a + 3 \end{pmatrix} \cdot \begin{pmatrix} a + 3 \\ -2 \end{pmatrix} = a^2 + 3a - 4a - 6 = 0$
$(a + 2)(a - 3) = 0 \Rightarrow a = -2, a = 3$
$a = -2 \Rightarrow \begin{pmatrix} -2 \\ -1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 \\ -2 \end{pmatrix}$
$a = 3 \Rightarrow \begin{pmatrix} 3 \\ 9 \end{pmatrix} \text{ and } \begin{pmatrix} 6 \\ -2 \end{pmatrix}$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ uses dot product to form equation</li> <li>✓ solves equation</li> <li>✓ states one pair of vectors</li> <li>✓ states both pairs of vectors</li> </ul>

Question 13


(8 marks)

- (a)  $A, B, C$  and  $D$  lie on a circle with diameter  $AC$  (diagram not to scale). Determine the size of  $\angle BDC$  when  $\angle BCA = 25^\circ$ . (2 marks)




Solution
$\angle BAC = 90 - 25 = 65$
$\angle BAD = \angle BAC = 65^\circ$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ uses angle in semi-circle for <math>\angle BAC</math></li> <li>✓ correct value</li> </ul>

- (b)  $K, L$  and  $M$  lie on a circle (diagram not to scale). Secant  $KN$  cuts the circle at  $L$  and  $JN$  is a tangent to the circle at  $M$ . Given that  $\angle LNM = 37^\circ$  and  $\angle LMN = 48^\circ$ , determine the size of  $\angle MKL$  and the size of  $\angle KMJ$ . (3 marks)



Solution
$\angle MKL = \angle LMN = 48^\circ$ (Alternate segments)
$\angle KLM = 37 + 48 = 85$ (Exterior angle)
$\angle KMJ = \angle KLM = 85^\circ$ (Alternate segments)
Specific behaviours
<ul style="list-style-type: none"> <li>✓ <math>\angle MKL</math></li> <li>✓ <math>\angle KLM</math></li> <li>✓ <math>\angle KMJ</math></li> </ul>

- (c)  $P, Q$  and  $R$  lie on a circle of radius 85 mm (diagram not to scale) and  $PQ = PR = 116$  mm. Determine the size of angle  $\angle QPR$ , to the nearest degree. (3 marks)



Solution
$116 \div 2 = 58$
$\theta = \cos^{-1} \frac{58}{85} = 47.0^\circ$
$\angle QPR = 2\theta = 94^\circ$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ completes diagram</li> <li>✓ uses trig ratio for half-angle</li> <li>✓ correct angle</li> </ul>

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

## Question 14

(9 marks)

The parts of this question refer to the word AERIFICATION. It has 5 different consonants and 7 vowels, some of which are repeated.

- (a) Determine the number of ways that 3 different consonants chosen from the letters of the word can be arranged in a row. (1 mark)

Solution
${}^5P_3 = 60$
Specific behaviours
✓ correct number

- (b) Determine the number of ways that all the letters of the word can be arranged in a row. (2 marks)

Solution
$\frac{12!}{3! \times 2!} = 39\,916\,800$
Specific behaviours
✓ attempts to account for repeated letters ✓ correct number

- (c) Determine the number of ways that all the letters of the word can be arranged in a row if the vowels must all be adjacent. (3 marks)

Solution
$\frac{(5 + 1)! \times 7!}{3! \times 2!} = 302\,400$
Specific behaviours
✓ counts vowels as single group ✓ counts ways to arrange vowels ✓ correct number

- (d) Determine how many 3 letter permutations (e.g. TFI, IRI, etc) can be made using the letters of the word. (3 marks)

Solution
All different: $9 \times 8 \times 7 = 504$
Two A's and one other: $3 \times 8 = 24$
Two I's and one other: $3 \times 8 = 24$
Three I's: 1
Total: $n = 504 + 24 + 24 + 1 = 553$
Specific behaviours
✓ attempts to consider separate cases ✓ correct number containing 2 A's and 2 I's ✓ correct total


See next page



Question 15

(8 marks)

- (a) In the diagram below (not drawn to scale)  $P, Q, R$  and  $S$  lie on the circle with centre  $O$ . Determine the size of angles  $\alpha, \beta$  and  $\gamma$  given that  $\angle PQR = 105^\circ$  and  $2\beta = 3\gamma$ . (4 marks)




Solution
$\alpha = 180 - 105 = 75^\circ$
$\beta + \gamma = 105$
$2\beta + 2\gamma = 210 \Rightarrow 5\gamma = 210 \Rightarrow \gamma = 42^\circ$
$\beta = 63^\circ$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ correct <math>\alpha</math></li> <li>✓ equation for <math>\beta + \gamma</math></li> <li>✓ correct <math>\gamma</math></li> <li>✓ correct <math>\beta</math></li> </ul>

- (b) Write the converse of the theorem that states the opposite angles of a cyclic quadrilateral are supplementary. (1 mark)

Solution
When opposite angles in a quadrilateral are supplementary, the quadrilateral is cyclic.
Specific behaviours
✓ correct statement

- (c) Prove by contradiction that the converse you wrote in (b) is true. Start by assuming that there is a quadrilateral that *does* have supplementary opposite angles but is *not* cyclic, such as  $STUV$  shown below. (3 marks)



Solution
From assumption, $\angle W = 180^\circ - \angle T$ .
But from regular theorem, $\angle V = 180^\circ - \angle T$ .
Hence $\angle W = \angle V$ , but this is impossible (as $SW$ and $SV$ would then be parallel and triangle $SVW$ would not exist). Thus, our original assumption must be wrong, and the converse must be true.
Specific behaviours
<ul style="list-style-type: none"> <li>✓ uses assumption</li> <li>✓ develops contradiction</li> <li>✓ explains contradiction and makes deduction</li> </ul>

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

## Question 16

(7 marks)

Three forces  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  act on a point in a plane.

The forces are  $\mathbf{a} = -44\mathbf{i} + 66\mathbf{j}$  N,  $\mathbf{b} = -12\mathbf{i} - 75\mathbf{j}$  N and  $\mathbf{c} = 180\mathbf{i} + 102\mathbf{j}$  N.

- (a) Determine the magnitude of the resultant force and the direction, to the nearest degree, that the resultant makes with the vector  $\mathbf{i}$ . (3 marks)

Solution
$\mathbf{r} = \begin{pmatrix} -44 \\ 66 \end{pmatrix} + \begin{pmatrix} -12 \\ -75 \end{pmatrix} + \begin{pmatrix} 180 \\ 102 \end{pmatrix} = \begin{pmatrix} 124 \\ 93 \end{pmatrix}$
$ \mathbf{r}  = 155 \text{ N}$
$\angle = 36.9 \approx 37^\circ$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ resultant</li> <li>✓ correct magnitude</li> <li>✓ correct angle</li> </ul>

When  $\lambda\mathbf{a} + \mu\mathbf{b} + \mathbf{c} = 0$ , the forces are in equilibrium.

- (b) Determine the values of the scalar constants  $\lambda$  and  $\mu$  for equilibrium to occur. (4 marks)

Solution
$\lambda \begin{pmatrix} -44 \\ 66 \end{pmatrix} + \mu \begin{pmatrix} -12 \\ -75 \end{pmatrix} + \begin{pmatrix} 180 \\ 102 \end{pmatrix} = 0$
$-44\lambda - 12\mu + 180 = 0$
$66\lambda - 75\mu + 102 = 0$
$\lambda = 3, \quad \mu = 4$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ equation using <math>\mathbf{i}</math>-coefficients</li> <li>✓ equation using <math>\mathbf{j}</math>-coefficients</li> <li>✓ solves for <math>\lambda</math></li> <li>✓ solves for <math>\mu</math></li> </ul>

Question 17

(8 marks)

- (a) A set of cards is numbered from 100 to 999. Determine the minimum number of cards that must be selected to ensure that at least 3 cards in the selection have the same last digit. Justify your answer using the pigeonhole principle. (3 marks)

<b>Solution</b>
Let pigeonholes be digits 0, 1, 2, ..., 9 and pigeons be the last digit of number on card.
Then fill all pigeonholes with 2 pigeons, a total of 20 pigeons.
The next pigeon will fill one of the pigeonholes with 3 pigeons, and so the minimum number is 21.
<b>Specific behaviours</b>
<ul style="list-style-type: none"> <li>✓ defines pigeons and pigeonholes</li> <li>✓ clear explanation</li> <li>✓ correct number</li> </ul>

- (b) Eight different books sit on a shelf, one of which has a hardcover and the rest softcovers. A student is told they can take away as many of them as they like but must not leave empty handed. Determine how many different selections can be made

- (i) of exactly 3 books. (1 mark)

<b>Solution</b>
$\binom{8}{3} = 56$
<b>Specific behaviours</b>
✓ correct number

- (ii) altogether. (2 marks)

<b>Solution</b>
Choose either 1, 2, ... up to all 8 books:
$\sum_{n=1}^8 \binom{8}{n} = 2^8 - 1 = 255$
<b>Specific behaviours</b>
<ul style="list-style-type: none"> <li>✓ uses property of Pascals triangle</li> <li>✓ correct number</li> </ul>

- (iii) that include the hardcover. (2 marks)

<b>Solution</b>
Choose hardcover and then 0, 1, ... up to 7 others:
$\binom{1}{1} \times \sum_{n=0}^7 \binom{7}{n} = 2^7 = 128$
<b>Specific behaviours</b>
<ul style="list-style-type: none"> <li>✓ indicates method</li> <li>✓ correct number</li> </ul>

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

## Question 18

(8 marks)

Relative to the origin,  $A$  and  $B$  have position vectors  $18\mathbf{i} + 18\mathbf{j}$  and  $21\mathbf{i} - 15\mathbf{j}$  respectively.

Particle  $P$  is initially at  $A$  and moves with a constant velocity of  $8\mathbf{i} - 15\mathbf{j} \text{ ms}^{-1}$ .

(a) Calculate

(i) the speed of  $P$ .

Solution	
$s = \sqrt{8^2 + (-15)^2} = 17 \text{ m/s}$	
Specific behaviours	
✓ correct speed	

(1 mark)

(ii) the position vector of  $P$  after 4 seconds.

Solution	
$\begin{pmatrix} 18 \\ 18 \end{pmatrix} + 4 \begin{pmatrix} 8 \\ -15 \end{pmatrix} = \begin{pmatrix} 50 \\ -42 \end{pmatrix}$	
Specific behaviours	
✓ correct position	

(1 mark)

(iii) the distance of  $P$  from  $B$  after 4 seconds.

Solution	
$\vec{PB} = \begin{pmatrix} 21 \\ -15 \end{pmatrix} - \begin{pmatrix} 50 \\ -42 \end{pmatrix} = \begin{pmatrix} -29 \\ 27 \end{pmatrix}$	
$ \vec{PB}  = \sqrt{(-29)^2 + (27)^2} = \sqrt{1570} \approx 39.6 \text{ m}$	
Specific behaviours	
✓ vector $\vec{PB}$	
✓ correct distance	

(2 marks)

(b) Determine how long after leaving  $A$  that  $P$  is 157 m from  $B$ .

(4 marks)

Solution	
$\vec{OP} = \begin{pmatrix} 18 \\ 18 \end{pmatrix} + t \begin{pmatrix} 8 \\ -15 \end{pmatrix}$	
$\vec{PB} = \begin{pmatrix} 21 \\ -15 \end{pmatrix} - \begin{pmatrix} 18 + 8t \\ 18 - 15t \end{pmatrix}$	
$ \vec{PB} ^2 = (3 - 8t)^2 + (-33 + 15t)^2 = 157^2$	
$t = 11$	
Specific behaviours	
✓ expression for $\vec{OP}$	
✓ expression for $\vec{PB}$	
✓ equation using distance	
✓ correct time	

Question 19

(7 marks)

$ABCD$  is a trapezium with  $\overrightarrow{AB}$  parallel and in the same direction to  $\overrightarrow{DC}$ .

(a) Sketch a labelled diagram of  $ABCD$ .

(1 mark)

Solution
Specific behaviours
✓ correct diagram

(b) Show that  $\overrightarrow{AC} + \overrightarrow{DB} = \overrightarrow{AB} + \overrightarrow{DC}$ .

(2 marks)

Solution
$\begin{aligned} \overrightarrow{AC} + \overrightarrow{DB} &= (\overrightarrow{AB} + \overrightarrow{BC}) + (\overrightarrow{DA} + \overrightarrow{AB}) \\ &= \overrightarrow{AB} + (\overrightarrow{DA} + \overrightarrow{AB} + \overrightarrow{BC}) \\ &= \overrightarrow{AB} + \overrightarrow{DC} \end{aligned}$
Specific behaviours
✓ splits $\overrightarrow{AC}$ and $\overrightarrow{DB}$ ✓ groups vectors that make $\overrightarrow{DC}$

(c)  $M$  lies on  $AC$  and  $N$  lies on  $BD$  so that  $AM:MC = BN:ND = 2:1$ . Use a vector method to prove that  $ABNM$  is a trapezium.

(4 marks)

Solution
$\overrightarrow{AM} = \frac{2}{3}\overrightarrow{AC}, \quad \overrightarrow{AN} = \overrightarrow{AB} + \frac{2}{3}\overrightarrow{BD}$
$\begin{aligned} \overrightarrow{NM} &= \overrightarrow{AM} - \overrightarrow{AN} \\ &= \frac{2}{3}\overrightarrow{AC} - \left(\overrightarrow{AB} + \frac{2}{3}\overrightarrow{BD}\right) \\ &= \frac{2}{3}(\overrightarrow{AC} - \overrightarrow{BD}) - \overrightarrow{AB} \\ &= \frac{2}{3}(\overrightarrow{AC} + \overrightarrow{DB}) - \overrightarrow{AB} \\ &= \frac{2}{3}(\overrightarrow{AB} + \overrightarrow{DC}) - \overrightarrow{AB} \text{ [from (b)]} \end{aligned}$
$\text{But } \overrightarrow{DC} = k\overrightarrow{AB}$
$\begin{aligned} \overrightarrow{NM} &= \frac{2}{3}(\overrightarrow{AB} + k\overrightarrow{AB}) - \overrightarrow{AB} \\ &= \left(\frac{2k-1}{3}\right)\overrightarrow{AB} \Rightarrow ABNM \text{ is trapezium} \end{aligned}$
Specific behaviours
✓ vectors for $M$ and $N$ ✓ obtains $\overrightarrow{NM}$ without $M$ and $N$ ✓ obtains $\overrightarrow{NM}$ in terms of $\overrightarrow{AB}, \overrightarrow{DC}$ ✓ obtains $\overrightarrow{NM}$ in terms of $\overrightarrow{AB}$

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

## Question 20

(7 marks)

Farm  $A$  lies 95 km away from farm  $B$  on a bearing of  $062^\circ$ . A helicopter leaves farm  $A$  at 7:30 am to fly to farm  $B$ . The helicopter can maintain a speed of  $145 \text{ kmh}^{-1}$  and there is a steady wind of  $35 \text{ kmh}^{-1}$  blowing from the north.

Determine the bearing that the helicopter should steer and the time of its arrival at farm  $B$ , to the nearest minute.

Solution
$\frac{\sin 62}{145t} = \frac{\sin \alpha}{35t}$
$\alpha = 12.3^\circ$
Bearing: $180 + 62 + 12.3 = 254.3^\circ$
$180 - 62 - 12.3 = 105.7$
$\frac{\sin 62}{145t} = \frac{\sin 105.7}{95}$
$t = 0.601 \text{ h}$ $= 36 \text{ m}$
Arrive at 8:06 am
Specific behaviours
<ul style="list-style-type: none"> <li>✓ diagram showing vectors and resultant</li> <li>✓ equation using sin rule for <math>\alpha</math></li> <li>✓ value of <math>\alpha</math></li> <li>✓ correct bearing</li> <li>✓ equation using sin rule for <math>t</math></li> <li>✓ value of <math>t</math></li> <li>✓ correct arrival time</li> </ul>

**Question 21**

**(8 marks)**

Determine how many of the integers between 1 and 340 inclusive are

(a) divisible by 6.

(1 mark)

Solution
$[340 \div 6] = 56$ $n = 56$
Specific behaviours
✓ correct number

(b) divisible by 6 or 7.

(3 marks)

Solution
LCM: $(6,7) = 42$ ;  $[340 \div 7] = 48$ $[340 \div 42] = 8$  $n = 56 + 48 - 8 = 96$
Specific behaviours
✓ number divisible by 42 ✓ indicates use of inclusion-exclusion ✓ correct number

(c) divisible by 6 or 7 but not both.

(1 mark)

Solution
$n = 96 - 8 = 88$
Specific behaviours
✓ correct number

(d) divisible by 6 or 7 but not 4.

(3 marks)

Solution
LCM's: $(6, 4) = 12$ ; $(7, 4) = 28$ ; $(4, 6, 7) = 84$  $[340 \div 12] = 28$ $[340 \div 28] = 12$ $[340 \div 84] = 4$  $n = 96 - 28 - 12 + 4 = 60$
Specific behaviours
✓ divisible by 12, 28 ✓ divisible by 84 ✓ correct number

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Supplementary page

Question number: \_\_\_\_\_

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF



Supplementary page

Question number: \_\_\_\_\_

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Supplementary page

Question number: \_\_\_\_\_

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Supplementary page

Question number: \_\_\_\_\_

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

