

SCHOOL

Trial WACE Examination, 2012

Question/Answer Booklet

**MATHEMATICS
SPECIALIST 3A/3B**
Section One:
Calculator-free

SOLUTIONS

Student Number: In figures

--	--	--	--	--	--	--	--

In words

Your name

MARTIN KEY

Time allowed for this section

Reading time before commencing work: five minutes

Working time for this section: fifty minutes

Materials required/recommended for this section

To be provided by the supervisor

This Question/Answer Booklet

Formula Sheet

To be provided by the candidate

Standard items: pens, pencils, pencil sharpener, eraser, correction fluid/tape, ruler, highlighters

Special items: nil

Important note to candidates

No other items may be used in this section of the examination. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of exam
Section One: Calculator-free	7	7	50	50	33
Section Two: Calculator-assumed	13	13	100	100	67
Total				150	100

Instructions to candidates

- The rules for the conduct of Western Australian external examinations are detailed in the *Year 12 Information Handbook 2012*. Sitting this examination implies that you agree to abide by these rules.
- Write your answers in the spaces provided in this Question/Answer Booklet. Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.
 - Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
 - Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.
- Show all your working clearly.** Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat an answer to any question, ensure that you cancel the answer you do not wish to have marked.
- It is recommended that you **do not use pencil**, except in diagrams.

Section One: Calculator-free

(50 Marks)

This section has **seven (7)** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time for this section is 50 minutes.

Question 1

(6 marks)

The complex numbers z and w are given by $z = 8 - 6i$ and $w = 3 + ai$.

(a) Determine $z \times \bar{z}$.

(2 marks)

$$\begin{aligned} z \times \bar{z} &= (8 - 6i)(8 + 6i) \checkmark \\ &= 64 + 36 \\ &= 100 \checkmark \end{aligned}$$

(b) Find $\text{Im}(w^2)$ if $a = 5$.

(2 marks)

$$\begin{aligned} w^2 &= (3 + 5i)(3 + 5i) \checkmark \\ &= 9 - 25 + 30i \checkmark \\ \text{Im}(w^2) &= 30 \checkmark \end{aligned}$$

(c) The value of a if $w^2 = z$.

(2 marks)

$$\begin{aligned} z &= 8 - 6i \\ w^2 &= (3 + ai)(3 + ai) \checkmark \\ &= 9 - a^2 + 6ai \checkmark \\ \text{From equating imaginary parts} \\ 6a &= -6 \\ a &= -1 \checkmark \end{aligned}$$

Question 2

(4 marks)

Determine the values of a and b if the function

$$f(x) = \begin{cases} a + \frac{4}{x} & 0 < x < 2 \\ bx^2 - 2 & x \geq 2 \end{cases}$$

is both continuous and differentiable at $x = 2$.

Continuous:

$$a + \frac{4}{2} = b \times 2^2 - 2 \Rightarrow a = 4b - 4 \quad \checkmark$$

Differentiable:

$$f'(x) = \begin{cases} -\frac{4}{x^2} & \checkmark \\ 2bx & \end{cases}$$

$$-\frac{4}{2^2} = 2 \times b \times 2 \Rightarrow b = -\frac{1}{4} \quad \checkmark$$

$$a = 4 \times -\frac{1}{4} - 4 = -5$$

$$\checkmark \quad a = -5, b = -\frac{1}{4}$$

Question 3

(9 marks)

(a) Solve $\log_x 64 = 3$.

(1 mark)

$$x^3 = 64 \Rightarrow x = 4 \quad \checkmark$$

(b) Rearrange the equation $11^{2x+3} = 33$ for x .

(2 marks)

$$\begin{aligned} (2x + 3) \log 11 &= \log 33 \quad \checkmark \\ 2x + 3 &= \frac{\log 33}{\log 11} \\ x &= \frac{\log 33}{2 \log 11} - \frac{3}{2} \quad \checkmark \end{aligned}$$

(c) Solve $2(\log_3 x + 1) = \log_3 25$.

(3 marks)

$$\begin{aligned} \log_3 x + 1 &= (0.5) \log_3 25 \quad \checkmark \\ \log_3 x + \log_3 3 &= \log_3 25^{0.5} \\ \log_3 3x &= \log_3 5 \quad \checkmark \\ x &= \frac{5}{3} \quad \checkmark \end{aligned}$$

(d) If $a = \log_5 4$ and $b = \log_5 8$, express the following in terms of a and b :

(i) $\log_5 32$

(1 mark)

$$\begin{aligned} \log_5 32 &= \log_5 (4 \times 8) \\ &= \log_5 4 + \log_5 8 \\ &= a + b \quad \checkmark \end{aligned}$$

(ii) $\log_5 400$


(2 marks)

$$\begin{aligned} \log_5 400 &= \log_5 (5 \times 4)^2 \quad \checkmark \\ &= 2(\log_5 5 + \log_5 4) \\ &= 2(1 + a) \quad \checkmark \end{aligned}$$

Question 4

(11 marks)

(a) The graph of the trigonometric function $y = f(x)$ is shown.



(i) State the amplitude of $f(x)$. (1 mark)

✓


(ii) State the period of $f(x)$. (1 mark)

✓

(iii) State the equation of the trigonometric function $f(x)$. (2 marks)

✓✓

- (b) The graphs of $g(x) = \frac{a}{b-x}$ and $h(x) = 4^{cx} + d$, where a, b, c and d are all non-zero constants are shown below, intersecting at $(-2, 3)$.



- (i) Explain why $g(x)$ cannot pass through the origin. (2 marks)

Because $g(0) = \frac{a}{b}$, but a is non-zero and so $g(0) \neq 0$.



- (ii) Find the values of a, b, c and d . (5 marks)

$g(x)$:
 $x=0 \Rightarrow 6 = \frac{a}{b} \Rightarrow a=6b$ ✓
 $x=-2 \Rightarrow 3 = \frac{a}{b+2}$ ✓
 $3b+6=6b$
 $b=2$ ✓
 $a=12$

 $h(x)$:
 $x=0 \Rightarrow 0=1+d \Rightarrow d=-1$ ✓
 $x=-2 \Rightarrow 3=4^{-2c}-1$
 $4=4^{-2c}$
 $-2c=1$ ✓
 $c=-0.5$ ✓

 $a=12, b=2, c=-0.5, d=-1$

Question 5

(8 marks)

Differentiate the following, simplifying where possible.

(a) $y = \frac{\log_e(4 - x^3)}{12}$

(2 marks)

$$\frac{dy}{dx} = \frac{1}{12} \times \frac{-3x^2}{4 - x^3}$$

$$= \frac{x^2}{4(x^3 - 4)}$$

(b) $y = \frac{3x}{2x+1} - \frac{1}{2x+1}$

(3 marks)

$$y = \frac{3x - 1}{2x + 1}$$

$$\frac{dy}{dx} = \frac{3(2x + 1) - 2(3x - 1)}{(2x + 1)^2}$$

$$= \frac{6x + 3 - 6x + 2}{(2x + 1)^2}$$

$$= \frac{5}{(2x + 1)^2}$$

(c) $y = (e^{\sqrt{x}} + 5)^4$

(3 marks)


$$\frac{dy}{dx} = 4 \times \left(\frac{1}{2\sqrt{x}} e^{\sqrt{x}} \right) (e^{\sqrt{x}} + 5)^3$$

$$= \frac{2e^{\sqrt{x}}}{\sqrt{x}} (e^{\sqrt{x}} + 5)^3$$

Question 6

(4 marks)

The graph of $y = f(x)$ is shown.



(a) Draw the graph of $y = f(|x|)$ on the same axes.

✓ idea
 ✓ accuracy within (sketch) reason (2 marks)

(b) Briefly explain how to obtain the graph of $y = f(|x|)$ from $y = f(x)$.

(2 marks)

Reflect the part of $y = f(x)$ in the 1st and 4th quadrants in the y-axis. ✓

Question 7

(8 marks)

(a) Show that $\tan(\theta) + \tan(2\theta) = \frac{\tan^3(\theta) - 3\tan(\theta)}{\tan^2(\theta) - 1}$.

(3 marks)

$$\begin{aligned} \tan(\theta) + \tan(2\theta) &= \tan(\theta) + \frac{2\tan(\theta)}{1 - \tan^2(\theta)} \quad \checkmark \\ &= \frac{\tan(\theta)(1 - \tan^2(\theta)) + 2\tan(\theta)}{1 - \tan^2(\theta)} \quad \checkmark \\ &= \frac{3\tan(\theta) - \tan^3(\theta)}{1 - \tan^2(\theta)} \\ &= \frac{\tan^3(\theta) - 3\tan(\theta)}{\tan^2(\theta) - 1} \quad \checkmark \end{aligned}$$

(b) Solve the equation $\tan(\theta) + \tan(2\theta) = 0$ for $0 \leq \theta \leq \pi$.

(5 marks)

$$\begin{aligned} \frac{\tan^3(\theta) - 3\tan(\theta)}{\tan^2(\theta) - 1} &= 0 \quad \checkmark \\ \tan^3(\theta) - 3\tan(\theta) &= 0 \\ \tan(\theta)(\tan^2(\theta) - 3) &= 0 \quad \checkmark \\ \tan(\theta)(\tan(\theta) + \sqrt{3})(\tan(\theta) - \sqrt{3}) &= 0 \\ \tan(\theta) = 0 &\Rightarrow \theta = 0, \pi \quad \checkmark \\ \tan(\theta) + \sqrt{3} = 0 &\Rightarrow \theta = \frac{2\pi}{3} \quad \checkmark \\ \tan(\theta) - \sqrt{3} = 0 &\Rightarrow \theta = \frac{\pi}{3} \quad \checkmark \\ \text{Solutions are } \theta &= 0, \frac{\pi}{3}, \frac{2\pi}{3}, \pi \end{aligned}$$

Additional working space

Question number: _____