



Christ Church Grammar School

Semester Two Examination, 2016

Question/Answer Booklet

**MATHEMATICS
SPECIALIST
UNITS 3 AND 4**
Section One:
Calculator-free

SOLUTIONS

Student Number: In figures

--	--	--	--	--	--	--	--

In words

Your name

Time allowed for this section

Reading time before commencing work: five minutes
Working time for section: fifty minutes

Materials required/recommended for this section

To be provided by the supervisor

This Question/Answer Booklet
Formula Sheet

To be provided by the candidate

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: nil

Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of exam
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	12	12	100	97	65
Total				149	100

Instructions to candidates

- The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
- Write your answers in this Question/Answer Booklet.
- You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
- Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.
 - Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
 - Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question that you are continuing to answer at the top of the page.
- Show all your working clearly.** Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
- It is recommended that you **do not use pencil**, except in diagrams.
- The Formula Sheet is **not** to be handed in with your Question/Booklet.

Section One: Calculator-free**35% (52 Marks)**

This section has **eight (8)** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time for this section is 50 minutes.

Question 1**(5 marks)**

The polynomial $h(z) = z^4 - 6z^3 + 3az^2 - 30z + 10a$, where a is a real constant, has a zero of $3 - i$. Determine the value of a and all other zeros of $h(z)$.

Solution
$h(z) = (z - 3 + i)(z - 3 - i) \times f(z)$
$z^4 - 6z^3 + 3az^2 - 30z + 10a = (z^2 - 6z + 10) \times f(z)$
<p>By inspection, $f(z) = (z^2 + bz + a)$</p>
$z^4 - 6z^3 + 3az^2 - 30z + 10a = (z^2 - 6z + 10)(z^2 + bz + a)$
<p>Comparing z^3 coefficients, $-6 = -6 + b \Rightarrow b = 0$</p>
<p>Comparing z^2 coefficients, $3a = 10 + a \Rightarrow a = 5$</p>
<p>Hence $h(z) = (z^2 - 6z + 10)(z^2 + 5)$</p>
<p>Other zeroes of $h(z)$ are $3 + i$, $\sqrt{5}i$ and $-\sqrt{5}i$.</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ deduces another zero of $3 + i$ ✓ uses two zeroes to obtain quadratic factor with real coefficients ✓ deduces other factor must be of form $z^2 + a$ ✓ determines value of a ✓ lists required zeroes

Question 2

(8 marks)

Two functions are defined by $f(x) = \sqrt{3x-1}$ and $g(x) = \frac{1}{x}$.


- (a) Determine the composite function $f(g(x))$ and the domain over which it is defined.

(3 marks)

Solution
$f(g(x)) = \sqrt{\frac{3}{x} - 1}$
Domain: $\frac{1}{x} \geq \frac{1}{3} \Rightarrow 0 < x \leq 3$.
Specific behaviours
<ul style="list-style-type: none"> ✓ writes composite function ✓ states upper limit of domain as $x \leq 3$ ✓ states lower limit of domain as $x > 0$

- (b) Sketch the graph of $y = g(g(x))$ on the axes below.


(2 marks)



Solution
$g(g(x)) = x, x \neq 0$
Specific behaviours
<ul style="list-style-type: none"> ✓ sketches graph of $y = x$ ✓ indicates discontinuity at $x = 0$

- (c) The graph of $y = f(x)$ is shown below, passing through point P with coordinates $(2.62, 2.62)$. Determine $f^{-1}(x)$, the inverse of $f(x)$, and sketch the graph of $y = f^{-1}(x)$ on the same axes.

(3 marks)



Solution
$x = \sqrt{3y-1} \Rightarrow y = f^{-1}(x) = \frac{x^2+1}{3}$
Specific behaviours
<ul style="list-style-type: none"> ✓ determines rule for inverse ✓ sketches inverse using reflection in $y = x$ ✓ inverse starts at $(0, \frac{1}{3})$ and passes thru P

Question 3

(5 marks)

An object, initially at rest, is dropped from the top of tall building so that after t seconds it has velocity v meters per second.

The air resistance encountered by the object is proportional to its velocity, so that the velocity satisfies the equation $\frac{dv}{dt} = 10 - kv$, where k is a constant.

(a) Express the velocity of the object in terms of t and k .

(4 marks)

Solution
$\int \frac{dv}{10 - kv} = \int dt$
$-\frac{1}{k} \ln 10 - kv = t + c$
$\ln 10 - kv = -kt - kc$
$10 - kv = ae^{-kt}$
$t = 0, v = 0 \Rightarrow a = 10$
$v = \frac{10 - 10e^{-kt}}{k}$
Specific behaviours
<ul style="list-style-type: none"> ✓ separates variables ✓ integrates both sides ✓ evaluates constant a ✓ expresses velocity as required

(b) Sensors on the object indicate that its velocity will never exceed 55 metres per second. Determine the value of the constant k .

(1 mark)

Solution
$\text{As } t \rightarrow \infty, 55 = \frac{10}{k} \Rightarrow k = \frac{10}{55} = \frac{2}{11}$
Specific behaviours
<ul style="list-style-type: none"> ✓ evaluates constant

Question 4

(5 marks)

Let $v = \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i$.

(a) Express v in polar form.

(2 marks)

Solution
$ v = 1, \arg v = -\frac{\pi}{4}, v = \text{cis}\left(-\frac{\pi}{4}\right)$
Specific behaviours
<ul style="list-style-type: none"> ✓ determines correct modulus ✓ determines correct argument


(b) Show that $v^4 = -1$.

(1 mark)

Solution
$v^4 = \left(\text{cis}\left(-\frac{\pi}{4}\right)\right)^4 = \text{cis}(-\pi) = -1$
Specific behaviours
✓ states v^4 in polar form

(c) Plot the roots of $z^4 + 1 = 0$ on the following Argand diagram.

(2 marks)



Solution
See diagram
Specific behaviours
<ul style="list-style-type: none"> ✓ adds scale and plots $v (= z_1)$ ✓ plots other three roots

Question 5

(8 marks)

- (a) Using partial fractions, or otherwise, determine $\int \frac{x-19}{(x+1)(x-4)} dx$.

(4 marks)

Solution
$A(x - 4) + B(x + 1) = x - 19 \Rightarrow A + B = 1, B - 4A = -19$
Solving gives $A = 4, B = -3$
$\int \frac{x - 19}{(x + 1)(x - 4)} dx = \int \frac{4}{x + 1} - \frac{3}{x - 4} dx$ $= 4 \ln x + 1 - 3 \ln x - 4 + c$
Specific behaviours
<ul style="list-style-type: none"> ✓ writes equations for A and B ✓ determines A and B ✓ integrates both fractions correctly ✓ includes constant of integration

- (b) Use the substitution $u = \sin x$ to evaluate $\int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{\cos x}{\sqrt{\sin x}} dx$.





(4 marks)

Solution
$du = \cos x dx, x = \frac{\pi}{2} \Rightarrow u = 1, x = \frac{\pi}{6} \Rightarrow u = \frac{1}{2}$
$\int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{\cos x}{\sqrt{\sin x}} dx = \int_{\frac{1}{2}}^1 \frac{1}{\sqrt{u}} du$ $= [2\sqrt{u}]_{\frac{1}{2}}^1$ $= 2 - \sqrt{2}$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses substitution correctly ✓ determines new bounds of integration ✓ integrates correctly ✓ evaluates integral

Question 6

(5 marks)

The differential equation $y' = \frac{2x}{y}$ is shown in just one of the four slope fields below.



Solution	
	See bottom left-hand slope field
Specific behaviours	
	<ul style="list-style-type: none"> ✓ chooses correct slope field ✓ uses slopes to sketch solution curve right of pt. ✓ uses slopes to sketch solution curve left of pt.

(a) On the slope field for $y' = \frac{2x}{y}$, sketch the solution of the differential equation that passes through the point (2, 4). (3 marks)

(b) Another solution to the differential equation passes through the point (6, -3). Use the incremental formula $\delta y \approx \frac{dy}{dx} \times \delta x$, with $\delta x = \frac{1}{10}$, to estimate the y-coordinate of this curve when $x = 6.1$. (2 marks)

Solution	
	$\delta y \approx \frac{2(6)}{-3} \times \frac{1}{10} = -0.4$ $y \approx -3 - 0.4 \approx -3.4$
Specific behaviours	
	<ul style="list-style-type: none"> ✓ calculates change in y ✓ calculates new y-coordinate

Question 7


(8 marks)

The function f is defined as $f(x) = \frac{x^2-1}{x^2+1}$.

- (a) Show that the **only** stationary point of the function occurs when $x = 0$. (2 marks)

Solution
$f'(x) = \frac{2x(x^2 + 1) - 2x(x^2 - 1)}{(x^2 + 1)^2} = \frac{4x}{(x^2 + 1)^2}$ <p>Hence $f'(x) = 0$ only when $4x = 0 \Rightarrow x = 0$.</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ differentiates function ✓ simplifies and makes conclusion

- (b) Sketch the graph of $y = f(x)$ on the axes below. (3 marks)



Solution
See graph
Specific behaviours
<ul style="list-style-type: none"> ✓ turning point ✓ roots ✓ approaches asymptote from below

- (c) Using your graph, or otherwise, determine all solutions to

- (i) $f(x) = |f(x)|$. (1 mark)

Solution
$x \leq -1 \cup x \geq 1$
Specific behaviours
✓ states both inequalities

- (ii) $f(x) = f(|x|)$. (1 mark)

Solution
$x \in \mathbb{R}$
Specific behaviours
✓ states all reals


- (iii) $f(x) = \frac{1}{f(x)}$. (1 mark)

Solution
$x = 0$
Specific behaviours
✓ states solution

Question 8

(8 marks)

A region is bounded by $x = 0$, $y = 0$ and $y = \ln(x + 3)$ as shown in the graph below.



- (a) Show that the area of the region is given by $\int_0^{\ln 3} (3 - e^y) dy$. (3 marks)
(Do not evaluate the integral).

Solution	
$y = \ln(x + 3) \Rightarrow x = e^y - 3, \quad x = 0 \Rightarrow y = \ln 3$	
Region is left of y -axis, so integral will be negative - need to multiply by -1 for area	
$A = - \int_0^{\ln 3} (e^y - 3) dy = \int_0^{\ln 3} (3 - e^y) dy$	
Specific behaviours	
<ul style="list-style-type: none"> ✓ expresses x in terms of y ✓ derives upper limit of integration ✓ explains need to multiply integral by -1, 	

- (b) Determine the volume of the solid obtained by revolving the region about the y -axis.

Solution	
$V = \pi \int_0^{\ln 3} (e^y - 3)^2 dy$	
$= \pi \int_0^{\ln 3} e^{2y} - 6e^y + 9 dy$	
$= \pi \left[\frac{e^{2y}}{2} - 6e^y + 9y \right]_0^{\ln 3}$	
$= \pi \left[\frac{9}{2} - 18 + 9 \ln 3 - \frac{1}{2} + 6 \right]$	
$= \pi(9 \ln 3 - 8)$	
Specific behaviours	
<ul style="list-style-type: none"> ✓ writes correct integral ✓ expands $(e^y - 3)^2$ ✓ integrates correctly ✓ substitutes upper and lower limits ✓ simplifies completely 	

(3 marks)

Additional working space

Question number: _____

© 2016 WA Exam Papers. Christ Church Grammar School has a non-exclusive licence to copy and communicate this paper for non-commercial, educational use within the school. No other copying, communication or use is permitted without the express written permission of WA Exam Papers.



Christ Church Grammar School

Semester Two Examination, 2016

Question/Answer Booklet

**MATHEMATICS
SPECIALIST
UNITS 3 AND 4**
Section Two:
Calculator-assumed

SOLUTIONS

Student Number: In figures

--	--	--	--	--	--	--	--

In words

Your name

Time allowed for this section

Reading time before commencing work: ten minutes
Working time for section: one hundred minutes

Materials required/recommended for this section

To be provided by the supervisor

This Question/Answer Booklet
Formula Sheet (retained from Section One)

To be provided by the candidate

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: drawing instruments, templates, notes on two unfolded sheets of A4 paper, and up to three calculators approved for use in the WACE examinations

Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of exam
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	12	12	100	97	65
Total				149	100

Instructions to candidates

1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer Booklet.
3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
4. Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.
 - Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
 - Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question that you are continuing to answer at the top of the page.
5. **Show all your working clearly.** Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you **do not use pencil**, except in diagrams.
7. The Formula Sheet is **not** to be handed in with your Question/Booklet.

Section Two: Calculator-assumed

65% (97 Marks)

This section has **twelve (12)** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time for this section is 100 minutes.

Question 9

(6 marks)

A system of equations is shown below.

$$\begin{aligned} x + 2y + 3z &= 1 \\ y + 3z &= -1 \\ -y + (a^2 - 4)z &= a + 2 \end{aligned}$$

(a) Determine the unique solution to the system when $a = 2$.

(2 marks)

Solution
$-y = 4 \Rightarrow y = -4$ $3z = -1 + 4 = 3 \Rightarrow z = 1$ $x - 8 + 3 = 1 \Rightarrow x = 6$ $x = 6, y = -4, z = 1$
Specific behaviours
✓ substitutes and solves for y ✓ solves for z and x

(b) Determine the value(s) of a so that the system

(i) has an infinite number of solutions.

(3 marks)

Solution
$(a^2 - 4 + 3)z = a + 2 - 1$ $(a + 1)(a - 1)z = a + 1$ Require $a = -1$ for an infinite number of solutions
Specific behaviours
✓ adds row2 and row 3 to eliminate y ✓ factorises $a^2 - 1$ ✓ states value of a

(ii) has no solutions.

(1 mark)

Solution
Require $a = 1$ for no solutions
Specific behaviours
✓ states value of a

Question 10

(8 marks)

The length of time, T months, that an athlete stays in an elite squad can be modelled by a normal distribution with population mean μ and population variance $\sigma^2 = 15$.

(a) An independent sample of five values of T is 7.7, 15.2, 3.9, 13.4 and 11.8 months.

- (i) Calculate the mean of this sample and state the distribution that a large number of such samples is expected to follow. (2 marks)

Solution
$\bar{x} = \frac{52}{5} = 10.4 \text{ months}$
<p>Sample means normally distributed with mean μ and variance $\sigma^2 = \frac{15}{5} = 3$ months</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ calculates sample mean ✓ states distribution with mean and variance

- (ii) Use this sample to construct a 90% confidence interval for μ , giving the bounds of the interval to two decimal places. (3 marks)

Solution
$z = 1.645$ for 90% CI
$10.4 \pm 1.645 \times \sqrt{3}$
<p>Interval is $7.55 < \mu < 13.25$ months</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ uses correct z-score ✓ calculates lower bound, rounding ✓ calculates upper bound, rounding

- (b) Determine the smallest number of values of T that would be required in a sample for the total width of a 95% confidence interval for μ to be less than 3 months. (3 marks)

Solution
$z = 1.96, \sigma = \sqrt{15}$
$n = \left(\frac{1.96 \times \sqrt{15}}{1.5} \right)^2 = 25.61$
<p>Smallest sample size is $n = 26$</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ shows z-score, variance and interval half-width used ✓ calculates n ✓ rounds n up

Question 11

(7 marks)

Plane p_1 has equation $3x + y + z = 6$ and line l has equation $\mathbf{r} = \mathbf{i} + \mathbf{j} + 2\mathbf{k} + t(\mathbf{i} - 2\mathbf{j} - \mathbf{k})$.

(a) Show that the line l lies in the plane p_1 .

(3 marks)

Solution
p_1 can be expressed as $\mathbf{r} \cdot \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix} = 6$ and l as $\mathbf{r} = \begin{bmatrix} 1+t \\ 1-2t \\ 2-t \end{bmatrix}$
Then $\begin{bmatrix} 1+t \\ 1-2t \\ 2-t \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix} = 6$
Which simplifies to $3 + 3t + 1 - 2t + 2 - t = 6$ and then to $6 = 6$
Hence line lies in plane.
Specific behaviours
<ul style="list-style-type: none"> ✓ writes plane in vector form ✓ substitutes line into plane equation ✓ simplifies and draws conclusion

(b) Another plane, p_2 , is perpendicular to plane p_1 , parallel to the line l and contains the point with position vector $\mathbf{i} - 3\mathbf{j} - \mathbf{k}$. Determine the equation of plane p_2 , giving your answer in the form $ax + by + cz = d$.

(4 marks)

Solution
p_2 must contain vectors $\begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix}$
$\mathbf{n} = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix} \times \begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \\ -7 \end{bmatrix}$
$\mathbf{r} \cdot \mathbf{n} = \begin{bmatrix} 1 \\ 4 \\ -7 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -3 \\ -1 \end{bmatrix} = -4$
Hence equation of p_2 is $\mathbf{r} \cdot \begin{bmatrix} 1 \\ 4 \\ -7 \end{bmatrix} = -4 \Rightarrow x + 4y - 7z = -4$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses two vectors in the plane ✓ calculates normal using cross product ✓ obtains vector equation of p_2 ✓ writes plane in required form

Question 12

(13 marks)

- (a) Show that the gradient of the curve $2x^2 + y^2 = 3xy$ at the point $(1, 2)$ is 2. (3 marks)

Solution
$2(2)x + 2yy' = 3y + 3xy'$ $x = 1, y = 2 \Rightarrow 4 + 4y' = 6 + 3y' \Rightarrow y' = 2$
Specific behaviours
<ul style="list-style-type: none"> ✓ correctly differentiates LHS ✓ correctly differentiates RHS ✓ substitutes x and y values and simplifies to show gradient

- (b) Another curve passing through the point $(-2, 10)$ has gradient given by $\frac{dy}{dx} = \frac{2xy}{1+x^2}$. Use a method involving separation of variables and integration to determine the equation of the curve. (4 marks)

Solution
$\int \frac{1}{y} dy = \int \frac{2x}{1+x^2} dx$ $\ln y = \ln(1+x^2) + c$ $y = k(1+x^2)$ $y = 10, x = -2 \Rightarrow k = 2$ $y = 2(1+x^2)$
Specific behaviours
<ul style="list-style-type: none"> ✓ separates variables ✓ integrates both sides ✓ eliminates natural logs ✓ determines constant and writes solution

(c) A particle is moving along the curve given by $y = \sqrt[3]{x}$, with one unit on both axes equal to one centimetre. When $x = 1$, the y -coordinate of the position of the particle is increasing at the rate of 2 centimetres per second.

(i) Show that the x -coordinate is increasing at 6 centimetres per second at this instant. (2 marks)

Solution
$\frac{dy}{dx} = \frac{1}{3}x^{-\frac{2}{3}}$ $\frac{dx}{dt} = \frac{dx}{dy} \times \frac{dy}{dt} = 3(1)^{\frac{2}{3}} \times 2 = 6 \text{ cm/s}$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses chain rule ✓ shows substitution of x-coordinate and given rate

(ii) Determine the exact rate at which the distance of the particle from the origin is changing at this instant. (4 marks)


Solution
$D = \sqrt{x^2 + y^2} = \sqrt{x^2 + x^{\frac{2}{3}}}$ $\frac{dD}{dx} = \frac{3x^{\frac{4}{3}} + 1}{3\sqrt[3]{x}\sqrt{x^2 + x^{\frac{2}{3}}}}$ $x = 1 \Rightarrow \frac{dD}{dx} = \frac{2\sqrt{2}}{3}$ $\frac{dD}{dt} = \frac{dD}{dx} \times \frac{dx}{dt} = \frac{2\sqrt{2}}{3} \times 6 = 4\sqrt{2} \text{ cm/s}$
Specific behaviours
<ul style="list-style-type: none"> ✓ determines expression for distance in terms of one variable ✓ differentiates expression wrt to variable ✓ substitutes, using chain rule ✓ simplifies in exact form

Question 13

(8 marks)

- (a) Sketch the graph of
- $y = \frac{2x+1}{x+1}$
- on the axes below.

(2 marks)



Solution
See graph
Specific behaviours
✓ axes intercepts and (1, 1)
✓ smooth curve

Simpson's rule is a formula used for numerical integration, the numerical approximation of definite integrals. When an interval $[a_0, a_n]$ is divided into an even number, n , of smaller intervals of equal width w , the bounds of these smaller intervals are denoted $a_0, a_1, a_2, \dots, a_{n-1}, a_n$. Simpson's rule can be expressed as follows:

$$\int_{a_0}^{a_n} f(x) dx = \frac{w}{3}(B + 2E + 4O)$$

where $B = f(a_0) + f(a_n)$, E is the sum of the values of $f(a_k)$ where k is even and O is the sum of the values of $f(a_k)$ where k is odd.

- (b) Use Simpson's rule with
- $n = 6$
- to evaluate an approximation for
- $\int_0^3 \frac{2x+1}{x+1} dx$
- , correct to four decimal places. (4 marks)

Solution
$a_0 = 1, a_1 = \frac{4}{3}, a_2 = \frac{3}{2}, a_3 = \frac{8}{5}, a_4 = \frac{5}{3}, a_5 = \frac{12}{7}, a_6 = \frac{7}{4}$
$B = \frac{11}{4}, E = \frac{19}{6}, O = \frac{488}{105}$
$\frac{1}{2} \left(\frac{11}{4} + 2 \times \frac{19}{6} + 4 \times \frac{488}{105} \right) = \frac{11623}{2520} \approx 4.6123$
Specific behaviours
✓ calculates ordinates
✓ calculates B, E and O
✓ evaluates using rule
✓ accurate to 4 dp

- (c) Determine the exact value of $\int_0^3 \frac{2x+1}{x+1} dx$ and hence calculate the percentage error of the approximation from (b). (2 marks)

Solution	
$\int_0^3 \frac{2x+1}{x+1} dx = 6 - 2 \ln 2.$	Hence $\frac{\left((6-2 \ln 2) - \frac{11623}{2520} \right)}{6-2 \ln 2} \times 100 \approx 0.03\%$ error.
Specific behaviours	
✓ calculates exact value	
✓ calculates % error	

Question 14

(7 marks)

- (a) The equation of a sphere with centre at $(2, -3, 1)$ is $x^2 + y^2 + z^2 = ax + by + cz - 2$.

Determine the values of a, b, c and the radius of the circle.

(3 marks)

Solution
Expanding $(x - 2)^2 + (y + 3)^2 + (z - 1)^2 = r^2$
Gives $x^2 + y^2 + z^2 - 4x + 6y - 2z + 14 = r^2$
Hence $a = 4, b = -6, c = 2$ and $r^2 - 14 = -2 \Rightarrow r = \sqrt{12} = 2\sqrt{3}$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses standard circle form and expands ✓ states values of a, b and c ✓ determines radius

- (b) Two particles, P and Q, leave their initial positions at the same time and travel with constant velocities shown in the table below.

Particle	Initial position	Velocity
P	$10\mathbf{i} - 5\mathbf{j} + 5\mathbf{k}$	$6\mathbf{i} + 2\mathbf{j} - 4\mathbf{k}$
Q	$28\mathbf{i} + 22\mathbf{j} - 31\mathbf{k}$	$2\mathbf{i} - 4\mathbf{j} + 4\mathbf{k}$

Show that the two particles collide, stating the position vector of the point of collision.

(4 marks)

Solution
$\mathbf{r}_P = \begin{bmatrix} 10 + 6t \\ -5 + 2t \\ 5 - 4t \end{bmatrix}, \quad \mathbf{r}_Q = \begin{bmatrix} 28 + 2t \\ 22 - 4t \\ -31 + 4t \end{bmatrix}$
Solving $10 + 6t = 28 + 2t \Rightarrow t = 4.5$.
$\mathbf{r}_P(4.5) = \begin{bmatrix} 37 \\ 4 \\ -13 \end{bmatrix}, \quad \mathbf{r}_Q(4.5) = \begin{bmatrix} 37 \\ 4 \\ -13 \end{bmatrix}$
Hence collide at $37\mathbf{i} + 4\mathbf{j} - 13\mathbf{k}$ when $t = 4.5$
Specific behaviours
<ul style="list-style-type: none"> ✓ states vector equations of both paths ✓ solves for t using one pair of coefficients ✓ substitutes to show both have same position ✓ states position vector of point of collision

Question 15

(8 marks)

- (a) Briefly describe a reason that a sample rather than a complete population may be used when carrying out a statistical investigation. (1 mark)

Solution
Cheaper, Quicker, Easier, etc.
Specific behaviours
✓ any legitimate reason

- (b) A researcher used government records to select a random sample of the ages of 114 men who had died recently in a town close to an industrial complex. The mean and standard deviation of the ages in the sample were 73.3 and 8.27 years respectively.

- (i) Explain why the sample standard deviation is a reasonable estimate for the population standard deviation in this case. (1 mark)

Solution
The sample size is large and well above 30.
Specific behaviours
✓ states large sample size

- (ii) Calculate a 98% confidence interval for the population mean and explain what the interval shows. (4 marks)

Solution
<p>z-score for 98% interval is 2.326</p> $73.3 \pm 2.326 \times \frac{8.27}{\sqrt{114}} = (71.5, 75.1)$ <p>We are 98% confident that the true mean lies between 71.5 and 75.1 years.</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ uses correct z-score ✓ demonstrates calculation used to calculate interval ✓ calculates correct interval ✓ explains interval


- (iii) The national average life-span of men was known to be 75.3 years. State with a reason what conclusion the researcher could draw from the confidence interval calculated in (ii) about the life-span of men in the town. (2 marks)

Solution
There is a significant difference in life-span of men in the town from those nationally, as the interval does not contain 75.3.
Specific behaviours
<ul style="list-style-type: none"> ✓ notes interval does not contain 75.3 ✓ uses 'difference' and 'lifespan' in explanation

Question 16

(8 marks)

- (a) On the Argand diagram below, clearly show the region that satisfies the complex inequalities given by $|z + 3 - 4i| \leq 5$ and $\frac{\pi}{2} \leq \arg z \leq \pi$. (4 marks)



Solution
See diagram, $ z - (-3 + 4i) \leq 5$
Specific behaviours
<ul style="list-style-type: none"> ✓ circle with centre $(-3, 4)$ and radius 5 ✓ circle intercepts at $(0, 0)$, $(0, 8)$, $(-6, 0)$ ✓ shades inside circle ✓ only shades in second quadrant


- (b) Determine all roots of the equation $z^5 = 16\sqrt{3} + 16i$, expressing them in the form $r \operatorname{cis} \theta$, where $r \geq 0$ and $-\pi \leq \theta \leq \pi$. (4 marks)

Solution
$z^5 = 32 \operatorname{cis} \frac{\pi}{6}$
$z_1 = 2 \operatorname{cis} \left(\frac{\pi}{30} \right)$
$z_2 = 2 \operatorname{cis} \left(\frac{13\pi}{30} \right)$
$z_3 = 2 \operatorname{cis} \left(\frac{25\pi}{30} \right)$
$z_4 = 2 \operatorname{cis} \left(\frac{-11\pi}{30} \right)$
$z_5 = 2 \operatorname{cis} \left(\frac{-23\pi}{30} \right)$
Specific behaviours
<ul style="list-style-type: none"> ✓ expresses z^5 in polar form ✓ determines first root ✓ determines other two roots with positive arguments ✓ determines other two roots with negative arguments

Question 17

(7 marks)

The graph of $y = f(x)$ is shown below, where $f(x) = 4\cos^4(2x)$ and A is the smallest root of $f(x)$, $x > 0$.



- (a) Show that $4\cos^4(2x) = \frac{3+4\cos(4x)+\cos(8x)}{2}$. (3 marks)

Solution
$4\cos^4(2x) = 4(\cos^2(2x))^2$ $= 4\left(\frac{1 + \cos 4x}{2}\right)^2$ $= 1 + 2\cos 4x + \cos^2 4x$ $= 1 + 2\cos 4x + \frac{1 + \cos 8x}{2}$ $= \frac{3 + 4\cos(4x) + \cos(8x)}{2}$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses double angle identity ✓ expands and uses identity again ✓ simplifies as required

- (b) Hence determine $\int \frac{3 + 4\cos(4x) + \cos(8x)}{2} dx$. (2 marks)

Solution
$\int \frac{3 + 4\cos(4x) + \cos(8x)}{2} dx = \frac{3x}{2} + \frac{1}{2}\sin 4x + \frac{1}{16}\sin 8x + c$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses result from (a) to integrate ✓ obtains correct result, including constant

- (c) Determine the exact volume of the solid generated when the region bounded by $y = f(x)$, $y = 0$, $x = 0$ and $x = A$ is rotated through 360° about the x -axis. (2 marks)

Solution
$\int_0^{\frac{\pi}{4}} \pi (4\cos^4(2x))^2 dx = \frac{35\pi^2}{32} \text{ cubic units}$
Specific behaviours
<ul style="list-style-type: none"> ✓ writes integral ✓ evaluates integral in exact form

Question 18

(10 marks)

- (a) A small object has initial position vector $\mathbf{r}(0) = \mathbf{i} + 3\mathbf{j} - \mathbf{k}$ metres and moves with velocity vector given by $\mathbf{v}(t) = 2t\mathbf{i} - 4t\mathbf{j} + 3\mathbf{k} \text{ ms}^{-1}$, where t is the time in seconds.
- (i) Show that the acceleration of the object is constant and state the magnitude of the acceleration. (2 marks)

Solution
$\mathbf{a}(t) = 2\mathbf{i} - 4\mathbf{j}$ - hence constant acceleration.
$ \mathbf{a} = \sqrt{2^2 + 4^2} = 2\sqrt{5} \text{ ms}^{-2}$
Specific behaviours
<ul style="list-style-type: none"> ✓ differentiates velocity ✓ evaluates magnitude

- (ii) Determine the position vector of the object after 2 seconds. (3 marks)

Solution
$\mathbf{r}(t) = t^2\mathbf{i} - 2t^2\mathbf{j} + 3t\mathbf{k} + \text{const}$
$\mathbf{r}(0) = \mathbf{i} + 3\mathbf{j} - \mathbf{k} \Rightarrow \mathbf{r}(t) = (t^2 + 1)\mathbf{i} + (3 - 2t^2)\mathbf{j} + (3t - 1)\mathbf{k}$
$\mathbf{r}(2) = (2^2 + 1)\mathbf{i} + (3 - 2(2)^2)\mathbf{j} + (3(2) - 1)\mathbf{k} = 5\mathbf{i} - 5\mathbf{j} + 5\mathbf{k}$
Specific behaviours
<ul style="list-style-type: none"> ✓ integrates velocity ✓ uses initial condition ✓ determines position vector

(b) Another small object has position vector given by $\mathbf{r}(t) = (1 + 2 \sec t)\mathbf{i} + (3 \tan t - 2)\mathbf{j}$ m, where t is the time in seconds.

(i) Determine the distance of the object from the origin when $t = \frac{\pi}{3}$. (2 marks)

Solution
$\left \mathbf{r}\left(\frac{\pi}{3}\right) \right = \left \left(1 + 2 \sec \frac{\pi}{3}\right)\mathbf{i} + \left(3 \tan \frac{\pi}{3} - 2\right)\mathbf{j} \right = 5\mathbf{i} + (3\sqrt{3} - 2)\mathbf{j} $ <p>Distance is 5.93 m (2 decimal places)</p>
Specific behaviours
<ul style="list-style-type: none"> ✓ substitutes $\frac{\pi}{3}$ and simplifies ✓ determines magnitude

(ii) Derive the Cartesian equation of the path of this object. (3 marks)

Solution
$x = 1 + 2 \sec t, \quad y = 3 \tan t - 2$ $\sec^2 t - \tan^2 t = 1 \Rightarrow \left(\frac{x-1}{2}\right)^2 - \left(\frac{y+2}{3}\right)^2 = 1$
Specific behaviours
<ul style="list-style-type: none"> ✓ equates x and y components ✓ uses trig identity ✓ rearranges and substitutes correctly

Question 19

(7 marks)

- (a) A particle undergoing simple harmonic motion with a period of 5 seconds is observed to move in a straight line, oscillating 3.6 m either side of a central position. Determine the speed of the particle when it is 3 m from the central position. (3 marks)

Solution
$\omega = \frac{2\pi}{5}$ $v^2 = \left(\frac{2\pi}{5}\right)^2 (3.6^2 - 3^2)$ $ v = 2.50 \text{ m/s}$
Specific behaviours
<ul style="list-style-type: none"> ✓ determines ω ✓ substitutes into velocity equation ✓ evaluates speed

- (b) Another particle moving in a straight line experiences an acceleration of $x + 2.5 \text{ ms}^{-2}$, where x is the position of the particle at time t seconds.

Given that when $x = 1$, the particle had a velocity of 2 ms^{-1} , determine the velocity of the particle when $x = 2$. (4 marks)

Solution
$\frac{1}{2}v^2 = \int x + 2.5 \, dx$ $v^2 = 2\left(\frac{x^2}{2} + 2.5x\right) + c$ $x = 1, v = 2 \Rightarrow c = -2$ $x = 2 \Rightarrow v^2 = 2\left(\frac{2^2}{2} + 2.5(2)\right) - 2 \Rightarrow v = \pm 2\sqrt{3}$
Specific behaviours
<ul style="list-style-type: none"> ✓ uses appropriate form of acceleration ✓ integrates ✓ evaluates constant ✓ states all possible values of v

Question 20

(8 marks)

The complex numbers w and z are given by $-\frac{1}{2} - \frac{\sqrt{3}}{2}i$ and $r(\cos \theta + i \sin \theta)$ respectively, where $r > 0$ and $-\frac{\pi}{3} < \theta < \frac{\pi}{3}$.

- (a) State, in terms of r and θ , the modulus and argument of wz and $\frac{z}{w}$. (3 marks)

Solution
$w = \text{cis} \frac{-2\pi}{3}, z = r \text{cis} \theta$
$wz = r \text{cis} \left(\theta - \frac{2\pi}{3} \right) \Rightarrow$ modulus is r and argument is $\theta - \frac{2\pi}{3}$
$\frac{z}{w} = r \text{cis} \left(\theta - \frac{-2\pi}{3} \right) \Rightarrow$ modulus is r and argument is $\theta + \frac{2\pi}{3}$
Specific behaviours
<ul style="list-style-type: none"> ✓ expresses w in cis form ✓ forms product and expresses modulus and argument ✓ forms quotient and expresses modulus and argument

- (b) Explain why the points represented by z, wz and $\frac{z}{w}$ in an Argand diagram are the vertices of an equilateral triangle. (2 marks)

Solution
The points are all at distance r from the origin and all subtend angles of $\frac{2\pi}{3}$ at the origin.
Specific behaviours
<ul style="list-style-type: none"> ✓ states equidistant from origin (or shows with diagram) ✓ states subtend equal angles at origin (or shows with diagram)

- (c) In an Argand diagram, one of the vertices of an equilateral triangle is represented by the complex number $5 - \sqrt{3}i$. If the other two vertices lie on a circle with centre at the origin, determine the complex numbers they represent in exact Cartesian form. (3 marks)

Solution
Let $z = 5 - \sqrt{3}i$ then other two vertices will be $wz = -4 - 2\sqrt{3}i$ and $\frac{z}{w} = -1 + 3\sqrt{3}i$
Specific behaviours
<ul style="list-style-type: none"> ✓ realises to use z ✓ determines first vertex ✓ determines second vertex

Additional working space

Question number: _____

Additional working space

Question number: _____

© 2016 WA Exam Papers. Christ Church Grammar School has a non-exclusive licence to copy and communicate this paper for non-commercial, educational use within the school. No other copying, communication or use is permitted without the express written permission of WA Exam Papers.