

Rossmoyne Senior High School

Semester Two Examination, 2017

Question/Answer booklet

MATHEMATICS SPECIALIST UNITS 3 AND 4 Section One: Calculator-free		SOLUTIONS
Student Number:	In figures	
	In words	
	Your name	

Time allowed for this section

Reading time before commencing work: Working time: five minutes fifty minutes

Materials required/recommended for this section

To be provided by the supervisor This Question/Answer booklet Formula sheet

To be provided by the candidate

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: nil

Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
				Total	100

Instructions to candidates

- 1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
- 2. Write your answers in this Question/Answer booklet.
- 3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
- 4. Additional working space pages at the end of this Question/Answer booklet are for planning or continuing an answer. If you use these pages, indicate at the original answer, the page number it is planned/continued on and write the question number being planned/continued on the additional working space page.
- 5. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
- 6. It is recommended that you do not use pencil, except in diagrams.
- 7. The Formula sheet is not to be handed in with your Question/Answer booklet.

2


Section One: Calculator-free

This section has **eight (8)** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time: 50 minutes.

Question 1

The graph of y = f(x) is shown below, where $f(x) = \log(17 - 4x)$.



(a) Explain why y is a function of x over the natural upinalit of f .	(a)	Explain why y is a function of x over the natural domain of f .	(1 mark)
---	-----	---	----------

Solution
A one-to-one relationship clearly exists.
Specific behaviours
✓ reasonable explanation

(b) Determine a function for the inverse of f.

Solution Let x = log(17 - 4y) $10^{x} = 17 - 4y$ $y = f^{-1}(x) = \frac{17 - 10^{x}}{4}$ Specific behaviours

✓ eliminates log

 \checkmark expresses as function of x

(c) On the axes above, sketch the graph of the inverse of f.

See next page

(2 marks)

(2 marks)

35% (52 Marks)

(5 marks)

(a) Given that z - 2 - i is a factor of $z^4 + az^2 + b$, determine the values of the real constants *a* and *b*. (4 marks)

SolutionLet z = 2 + i $z^2 = (2 + i)(2 + i) = 3 + 4i$ $z^4 = (3 + 4i)(3 + 4i) = -7 + 24i$ -7 + 24i + 3a + 4ai + b = 0Im: $24 + 4a = 0 \Rightarrow a = -6$ Re: $-7 - 18 + b = 0 \Rightarrow b = 25$ a = -6, b = 25Specific behaviours \checkmark identifies root and substitutes \checkmark determines z^4 correctly \checkmark equates Re and Im parts \checkmark solves for constants

(b) Solve the equation $z^4 + z^2 - 12 = 0$.

Solution $(z^2 + 4)(z^2 - 3) = 0$ $z = \pm 2i, \quad z = \pm \sqrt{3}$ Specific behaviours \checkmark factorises \checkmark Re solutions \checkmark Im solutions (3 marks)

Let $u = 1 + \sqrt{3}i$ and z = -1 - i.

(a) Determine the argument of u^4z .

Solution
$\arg u^{4} = 4 \times \frac{\pi}{3} = \frac{4\pi}{3}$ $\arg z = -\frac{3\pi}{4}$ $\arg(u^{4}z) = \frac{4\pi}{3} + \left(-\frac{3\pi}{4}\right) = \frac{7\pi}{12}$
Specific behaviours
\checkmark indicates $\arg u^4$ and $\arg z$
✓ indicates addition of arguments
✓ states required argument

(b) Determine the real part of $(iu)^6$.

Solution
$$iu = -\sqrt{3} + i$$
 $= 2 \operatorname{cis} \frac{5\pi}{6}$ $(iu)^6 = 2^6 \operatorname{cis} \pi$ Re part $= -2^6 = -64$ Specific behaviours \checkmark indicates iu \checkmark expresses iu in polar form \checkmark determines $(iu)^6$ \checkmark states real part

(4 marks)

SPECIALIST UNITS 3 AND 4

(3 marks)

CALCULATOR-FREE

Question 4

(a) Determine the slope of the graph of $2x^2 + y^2 = 3xy$ at the point (1, 2).

(7 marks) (3 marks)

Solution4x + 2yy' = 3y + 3xy'4 + 4y' = 6 + 3y'y' = 2Specific behaviours \checkmark differentiates LHS \checkmark differentiates RHS \checkmark substitutes and simplifies

(b) Use the substitution $u = 5x^2 - 4$ to express $\int_{1}^{2} \frac{10x}{\sqrt{5x^2 - 4}} dx$ in terms of u and hence evaluate the integral. (4 marks)

Solution

$$u = 5x^{2} - 4 \Rightarrow du = 10x \, dx$$

$$x = 1, u = 1; \ x = 2, u = 16$$

$$\int_{1}^{2} \frac{10x}{\sqrt{5x^{2} - 4}} \, dx = \int_{1}^{16} \frac{1}{\sqrt{u}} \, du$$

$$= \left[2\sqrt{u}\right]_{1}^{16} = 8 - 2 = 6$$
Specific behaviours
 \checkmark expresses integrand in terms of u
 \checkmark changes limits
 \checkmark antidifferentiates
 \checkmark evaluates

6

Two functions are defined by

$$f(x) = \frac{x+1}{1-x^2}$$
 and $g(x) = x-3$.

(a) Determine
$$g \circ f(-2)$$
.

	Solution	
$f(-2) = \frac{1}{3},$	$g\left(\frac{1}{3}\right) = -\frac{8}{3},$	$g \circ f(-2) = -\frac{8}{3}$
	Specific behavior	ours
✓ evaluates		

(b) Determine a simplified expression for $f \circ g(x)$ and state the domain and range for this composite function. (5 marks)


Solution
$$f(x) = \frac{1+x}{(1-x)(1+x)}$$
 $f \circ g(x) = \frac{1+x-3}{(1-x+3)(1+x-3)}$ $= \frac{1}{4-x}, x \neq 2$ $D_{fog} = \{x: x \neq 4, x \neq 2\}$ $R_{fog} = \{y: y \neq 0, y \neq \frac{1}{2}\}$ Specific behaviours \checkmark correct expression for composite \checkmark simplifies composite \checkmark correct domain \checkmark range: includes $y \neq 0$ \checkmark correct range

(6 marks)

7

(5 marks)


A plane passes through the points A(0, 1, 0), B(1, 3, -2) and C(-2, -1, 1). Determine the Cartesian equation of the plane.



(7 marks)


(a) Part of the graph with equation
$$y = \frac{-36}{x^2 - 9}$$
 is shown below.

Complete the sketch, clearly indicating the location of any asymptotes. (2 marks)



(b) Determine the area bounded by $y = \frac{-36}{x^2 - 9}$, the *x*-axis, the *y*-axis and the line x = -2.

(5 marks)



SPECIALIST UNITS 3 AND 4

Question 8

(8 marks)

(a) Determine all solutions to the complex equation $3^3 z^6 = \frac{1}{2}(-\sqrt{3}+i)$ in the form $r \operatorname{cis} \theta$ where r > 0 and $-\pi < \theta \le \pi$. (5 marks)

10

Solution
$3^3 z^6 = \operatorname{cis} \frac{5\pi}{6}$
$z^6 = \frac{1}{3^3} \operatorname{cis} \frac{5\pi}{6}$
$z = \frac{1}{\sqrt{3}} \operatorname{cis}\left(\frac{5\pi}{36} + \frac{k\pi}{3}\right), k = -3, -2, -1, 0, 1, 2$
$z_1 = \frac{1}{\sqrt{3}} \operatorname{cis} \frac{-31\pi}{36}$
$z_2 = \frac{1}{\sqrt{3}} \operatorname{cis} \frac{-19\pi}{36}$
$z_3 = \frac{1}{\sqrt{3}} \operatorname{cis} \frac{-7\pi}{36}$
$z_4 = \frac{1}{\sqrt{3}} \operatorname{cis} \frac{5\pi}{36}$
$z_5 = \frac{1}{\sqrt{3}} \operatorname{cis} \frac{17\pi}{36}$
$z_5 = \frac{1}{\sqrt{3}} \operatorname{cis} \frac{29\pi}{36}$
Specific behaviours
\checkmark expresses z^6 in polar form
✓ uses De Moivre's theorem
\checkmark expresses general solution in terms of k
✓ states one correct root
✓ states all six roots correctly

(b) If w is any complex fifth root of unity, simplify $(1 + w)(1 + w^2)(1 + w^3)$.

(3 marks)

Solution $w^5 = 1 \text{ and } 1 + w + w^2 + w^3 + w^4 = 0 \text{ hence}$ $(1+w)(1+w^2)(1+w^3) = (1+w+w^2+w^3)(1+w^3)$ $= 1+w+w^2+w^3+w^3+w^4+w^5+w^6$ $= w^3+1+w^6$ $= w^3+w+1$ Specific behaviours \checkmark uses fifth power of root \checkmark uses sum of roots \checkmark simplifies Additional working space

Question number: _____

© 2017 WA Exam Papers. Rossmoyne Senior High School has a non-exclusive licence to copy and communicate this document for non-commercial, educational use within the school. No other copying, communication or use is permitted without the express written permission of WA Exam Papers. SN085-104-2.