SCOTCH COLLEGE	Year 12 Specialist Mathematics	
X	TEST 3 – Integration	
	Part A – Calculator Free	
Name:	Teacher: Fugill Langley	(please circle)

Date: 6th April 2016

SECTION ONE: CALCULATOR FREE

TOTAL: 35 marks

EQUIPMENT: Pens, pencils, pencil sharpener, highlighter, eraser, ruler, SCSA formula sheet.

WORKING TIME: 35 minutes

SECTION TWO: CALCULATOR ASSUMED

TOTAL: 19 marks

EQUIPMENT: Pens, pencils, pencil sharpener, highlighter, eraser, ruler, drawing instruments, templates, up to 3 Calculators,

1 A4 page of notes (one side only), SCSA formula sheet.

WORKING TIME: 20 minutes

Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat an answer to any question, ensure that you cancel the answer you do not wish to have marked.

Question	Marks available	Marks awarded
1	10	
2	8	
3	4	
4	6	
5	7	
6	5	
7	6	
8	8	
Total	54	

Calculator-free

[35 marks]

This paper has Five (5) questions. Answer all questions. Write your answers in the spaces provided

Question 1 [10 marks]

Determine the following indefinite integrals.

a)
$$\int \frac{x^2 + 2x - 5}{x^2} dx$$

[3]

[3]

b) $\int \sin^3(2x) \cos(2x) \, dx$

c) $\int x\sqrt{x-1} dx$

[4]

Evaluate:

a)
$$\int_0^{\frac{\pi}{4}} \cos 2x \, dx$$


[3]

b)
$$\int_0^1 \frac{x-1}{x^2+3x+2} dx$$

[5]

Question 3 [4 marks]

The curves below are $y = \cos(2x)$ and $y = \cos^2 x$. Determine the area of the shaded region.




Question 4 [6 marks]

A glass is formed by rotating the function $f(x) = |x^2 - 1|$ about the y-axis as shown in the diagram.

a) Complete the middle line for the piecewise function for $f(x) = |x^2 - 1|$.

$$f(x) = \begin{cases} x^2 - 1 \ ; \ x \le -1 \\ x^2 - 1 \ ; \ x \ge 1 \end{cases}$$



[1]

[5]

b) Given the height of the glass is 8 cm determine the volume of the glass.

Question 5 [7 marks]

Find $\int_{1}^{\sqrt{3}} \sqrt{4-x^2} dx$ using the substitution $x = 2 \sin \theta$.

SCOTCH COLLEGE	Year 12 Specialist Mathematics	
	TEST 3 – Integration	
	Part B – Calculator Allowed	
Name:	Teacher: Fugill Langley (please circle)	

Calculator Allowed

20 minutes

[19 marks]

This paper has Three (3) questions. Answer all questions. Write your answers in the spaces provided

Question 6 [5 marks]

Determine the value of p in the system of linear equations below such that there is

a) no solution	x - 2y - 3z = 11	[3]
	2x - y + z = 5	[9]
	3x + 3y + pz = -6	

b) a unique solution

c) infinitely many solutions

[1]


[1]

Question 7 [6 marks]

The graph at the right show the curves $y = \cos x$ and $y = \sin x$.

a) Prove that the intersection of $y = \cos x$ and $y = \sin x$, is $x = \frac{\pi}{4}$

for the domain $0 \le x \le \frac{\pi}{2}$.



b) Determine the exact area of the region (shaded) which is bounded by the *y*-axis and the curves $y = \cos x$ and $y = \sin x$. [2]

c) Determine the volume of revolution obtained when this area is rotated about the *x*-axis. [3]

Question 8 [8 marks]

a) Use the identity $\cos 2\theta = 2\cos^2 \theta - 1$ to prove that $\cos\left(\frac{1}{2}x\right) = \sqrt{\frac{1+\cos x}{2}}, \ 0 \le x \le \pi.$ [2]

b) Find a similar expression for $\sin\left(\frac{1}{2}x\right)$, $0 \le x \le \pi$. [2]

c) Hence show that
$$\int_{0}^{\frac{\pi}{2}} (\sqrt{1 + \cos x} + \sqrt{1 - \cos x}) dx = 2\sqrt{2}$$
 [4]