 <p>PERTH MODERN SCHOOL Exceptional schooling. Exceptional students. Independent Public School</p>	<p>Year 12 Specialist TEST 1 Friday 9 February 2018 TIME: 5 mins reading 40 minutes working Classpads allowed! 37 marks 7 Questions</p>
--	--

Name: SOLUTIONS

Teacher: _____

Note: All part questions worth more than 2 marks require working to obtain full marks.

Some useful Formulae

Cartesian form	
$z = a + bi$	$\bar{z} = a - bi$
$\text{Mod}(z) = z = \sqrt{a^2 + b^2} = r$	$\text{Arg}(z) = \theta, \quad \tan \theta = \frac{b}{a}, \quad -\pi < \theta \leq \pi$
$ z_1 z_2 = z_1 z_2 $	$\left \frac{z_1}{z_2} \right = \frac{ z_1 }{ z_2 }$
$\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$	$\arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2)$
$z \bar{z} = z ^2$	$z^{-1} = \frac{1}{z} = \frac{\bar{z}}{ z ^2}$
$\overline{z_1 + z_2} = \bar{z}_1 + \bar{z}_2$	$\overline{z_1 z_2} = \bar{z}_1 \bar{z}_2$
Polar form	
$z = a + bi = r(\cos \theta + i \sin \theta) = r \text{cis } \theta$	$\bar{z} = r \text{cis } (-\theta)$
$z_1 z_2 = r_1 r_2 \text{cis}(\theta_1 + \theta_2)$	$\frac{z_1}{z_2} = \frac{r_1}{r_2} \text{cis}(\theta_1 - \theta_2)$
$\text{cis}(\theta_1 + \theta_2) = \text{cis } \theta_1 \text{cis } \theta_2$	$\text{cis}(-\theta) = \frac{1}{\text{cis } \theta}$
De Moivre's theorem	
$z^n = z ^n \text{cis}(n\theta)$	$(\text{cis } \theta)^n = \cos n\theta + i \sin n\theta$
$z^{\frac{1}{q}} = r^{\frac{1}{q}} \left(\cos \frac{\theta + 2\pi k}{q} + i \sin \frac{\theta + 2\pi k}{q} \right), \quad \text{for } k \text{ an integer}$	

$\cos^2 x + \sin^2 x = 1$	$1 + \tan^2 x = \sec^2 x$
$\cos(x \pm y) = \cos x \cos y \mp \sin x \sin y$	$\cos 2x = \cos^2 x - \sin^2 x$ $= 2 \cos^2 x - 1$ $= 1 - 2 \sin^2 x$
$\sin(x \pm y) = \sin x \cos y \pm \cos x \sin y$	$\sin 2x = 2 \sin x \cos x$
$\tan(x \pm y) = \frac{\tan x \pm \tan y}{1 \mp \tan x \tan y}$	$\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$
$\cos A \cos B = \frac{1}{2}(\cos(A - B) + \cos(A + B))$	$\sin A \cos B = \frac{1}{2}(\sin(A + B) + \sin(A - B))$
$\sin A \sin B = \frac{1}{2}(\cos(A - B) - \cos(A + B))$	$\cos A \sin B = \frac{1}{2}(\sin(A + B) - \sin(A - B))$

Note: All part questions worth more than 2 marks require working to obtain full marks.

Q1) (2, 2, 2, 2 & 1 = 9 marks)

If $w = 2 - 2i$ and $z = 9 - 5i$ determine exactly:

a) wz

$$8 - 28i$$

✓ Real term ✓ Imaginary

b) $\frac{w}{z}$

$$\frac{2-2i}{9-5i} \cdot \frac{9+5i}{9+5i} = \frac{28-8i}{9^2+25^2} = \frac{28-8i}{706}$$

✓ numerator
✓ denominator

c) $z\bar{w}$

~~$8 - 28i$~~ $28 + 8i$

✓ Real ✓ Imaginary

d) $w\bar{z}$

$28 - 8i$

✓ Real ✓ Imaginary

e) What do you notice about (c) and (d)?

conjugates


✓ mentions conjugates

Q2 (2 & 2 = 4 marks)

Express each of the following into Cartesian form, $a + bi$

a) $7\text{cis}\left(-\frac{2\pi}{3}\right) = 7\left(\cos\left(-\frac{2\pi}{3}\right) + i\sin\left(-\frac{2\pi}{3}\right)\right) = 7\left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = -\frac{7}{2} - \frac{7\sqrt{3}}{2}i$

✓ expands cis
✓ evaluates Re & Im parts




$$12\cos 30^\circ - 12\sin 30^\circ i = 6\sqrt{3} - 6i$$

✓ real part

✓ Imaginary part

Q4 (3 marks)

Determine z in polar form given that w and zw have been drawn below.



~~5 cis 30°~~ $r \text{ cis } \theta = 8 \text{ cis}(220^\circ)$

$r = \frac{8}{5}$ $\theta = 190^\circ$

$z = \frac{8}{5} \text{ cis } 190^\circ$ or $\frac{8}{5} \text{ cis }(-170^\circ)$

- ✓ uses additive property of cis
- ✓ determines r
- ✓ determines θ

Q5 (5, 3 & 3 = 11 marks)

a) Determine all the roots of the equation $z^5 = 1 - i$, expressing them all in polar form with $r \geq 0$ and $-\pi < \text{Arg}z \leq \pi$

$z^5 = \sqrt{2} \text{ cis}(-\frac{\pi}{4} + 2n\pi)$ $n=0, \pm 1, \pm 2, \dots$

$z = \sqrt[5]{2} \text{ cis}(-\frac{\pi}{20} + \frac{2n\pi}{5})$

$= 2^{\frac{1}{10}} \text{ cis}(-\frac{\pi}{20} + \frac{8n\pi}{20})$

$z_1 = 2^{\frac{1}{10}} \text{ cis}(-\frac{\pi}{20})$ ✓ uses $2^{\frac{1}{10}}$


$z_2 = 2^{\frac{1}{10}} \text{ cis}(\frac{7\pi}{20})$ ✓ identifies $-\frac{\pi}{20}$

$z_3 = 2^{\frac{1}{10}} \text{ cis}(-\frac{9\pi}{20})$ ✓ determines 5 different arguments

$z_4 = 2^{\frac{1}{10}} \text{ cis}(\frac{15\pi}{20})$ ✓ converts to principal Arg

$z_5 = 2^{\frac{1}{10}} \text{ cis}(-\frac{17\pi}{20})$ ✓ states all 5 roots

b) Plot the roots on the diagram below. (Note: each minor angle is $\frac{\pi}{20}$ radians.)




✓ shows scale ($r = 2^{\frac{1}{10}}$)

✓ Five equally spaced points

✓ all 5 pts have correct angle.

- c) The roots form the vertices of a pentagon. Determine the value for the perimeter of the pentagon to two decimal places.



$$\sin \frac{\pi}{5} = \frac{2^{1/10}}{x}$$

$$x = 2^{1/10} \sin \frac{\pi}{5}$$

✓ using correct angle.

$$\text{Perimeter} = 10 \left(2^{1/10} \sin \frac{\pi}{5} \right)$$

✓ solving opposite side of triangle of angle.

$$= 6.30 \text{ units}$$

✓ determines perimeter

Q6 (5 marks)

Determine, using de Moivre's theorem, an expression for $\sin 3\theta$ in terms of $\sin \theta$ only.

{Hint: start with $(\cos \theta + i \sin \theta)^3$ }

$$(\cos \theta + i \sin \theta)^3 = \text{cis } 3\theta$$

$$= \cos^3 \theta - 3 \cos \theta \sin^2 \theta - (\sin^3 \theta - 3 \cos^2 \theta \sin \theta) i$$

$$= \cos^3 \theta + i \sin^3 \theta$$

$$\sin 3\theta = \sin^3 \theta + 3 \cos^2 \theta \sin \theta$$

$$= \sin^3 \theta + 3(1 - \sin^2 \theta) \sin \theta$$

$$= \sin^3 \theta + 3 \sin \theta - 3 \sin^3 \theta$$

$$= 3 \sin \theta - 2 \sin^3 \theta$$

✓ equates $(\cos \theta + i \sin \theta)^3$ to $\text{cis } 3\theta$

✓ expands $(\cos \theta + i \sin \theta)^3$


✓ equates Im part to $\sin 3\theta$

✓ replaces $\cos^2 \theta$ with $1 - \sin^2 \theta$

✓ obtains final expression in terms of $\sin \theta$

Q7 (5 marks)

Consider the points A and B in the complex plane. The perpendicular bisector of the line AB is represented by $\text{Im}(z) = 3\text{Re}(z) - 1$



If point A is $5 + ci$ and point B is $d - 7i$ in the complex plane, determine the values of the constants c and d .

$$\text{Midpoint } AB = \left(\frac{5+d}{2}, \frac{c-7}{2} \right) \quad \frac{c-7}{2} = 3 \left(\frac{5+d}{2} \right) - 1$$

$$m_{AB} = \frac{c+7}{5-d} = -\frac{1}{3}$$

Use simultaneous: $c = -12\frac{1}{4}$

$$d = -10\frac{3}{4}$$

- ✓ determines midpoint in terms of c & d
- ✓ determines gradient in terms of c & d
- ✓ obtains one equation, c & d (ie midpoint into line eqn)
- ✓ obtains two equations, c & d (ie $m_1 \times m_2 = -1$)
- ✓ Solves for c & d .